

52
671
M251
BIRD

MALIMBUS

Journal of West African Ornithology
Revue d'Ornithologie de l'Ouest Africain

50 years
of
N.O.S./W.A.O.S.
1964-2014



VOLUME 36 Number 2
ISSN 0331-3689

September 2014

published by:
publiée par:

West African Ornithological Society
Société d'Ornithologie de l'Ouest Africain

West African Ornithological Society Société d'Ornithologie de l'Ouest Africain

Conseil:

Président: Dr Jean-Marc Thiollay
Vice-Président: Dr Roger Wilkinson
Membres du Conseil:

Nils Robin, P.W. Peter Browne

Secrétaire du Conseil: Dr Joost Brouwer
Trésorier et chargé des adhérents: Tim Dodman
Secrétaire chargé des Réunions: Dr Shiiwua Manu
Rédacteur en chef: Dr Alan Tye

Comité de Rédaction: Dr J. Brouwer, Dr R. Buij, Dr D.T. Cox, Dr O. Girard, Dr P.J. Jones, Dr O. Lachenaud, Dr M. Melo, N. Robin, Dr V. Salewski, J.-M. Thiollay, Dr T. Wacher, Dr R. Wilkinson.

Site internet: <http://malimbus.free.fr/> inclut le texte complet de tous les volumes du *Bulletin of the Nigerian Ornithologists' Society* et de *Malimbus* à l'exception de ceux des cinq dernières années, ainsi que les tables des matières, les résumés, les index consultables d'espèces et d'auteurs, et des listes pays avec, pour chacun des pays d'Afrique de l'Ouest, les références des mentions dans la revue. Webmestre: Peter Browne.

La correspondance doit être adressée comme suit:

— au Rédacteur en Chef, pour les articles à soumettre à *Malimbus* (Dr A. Tye, PO Box 31, Kalo Chorio Oreinis, 2616 Nicosia, Cyprus ; <alantye@gmail.com>);

— au Trésorier, pour les abonnements (Hundland, Papa Westray, Orkney KW17 2BU, U.K.; <tim@timdodman.co.uk>);

— au Secrétaire du Conseil, pour les demandes des Bourses de Recherches (Wildekamp 32, 6721 JD Bennekom, The Netherlands; <brouwereac@orange.nl>);

— au Secrétaire adjoint, pour la présence aux réunions ou des suggestions pour celles-ci (APLORI, c/o P. Hall, A.G. Leventis Ltd, West Africa House, Hanger Lane, Ealing, London W5 3QR, U.K.; <manushiiwua@yahoo.co.uk>)

— au Webmestre, Peter Browne, pour les questions du site web: <pbrowne@primus.ca>;

— au Président, pour les questions du politique de la Société (2 rue Rivière, F-10220 Rouilly Sacey, France; <jm.thiollay@wanadoo.fr>).

La Société tire son origine de la "Nigerian Ornithologists' Society", fondée en 1964. Son but est de promouvoir l'ornithologie ouest-africaine, principalement au moyen de sa revue *Malimbus* (anciennement *Bulletin of the Nigerian Ornithologists' Society*).

Les demandes d'adhésion sont les bienvenues. Les cotisations annuelles sont de £15 ou de €22 pour les Membres Ordinaires et de £35 (€44) pour les Sociétés (les cotisations peuvent être payées en £ sterling au Trésorier ou en Euro à M. ou Mme N. Robin, 35 rue Bonaparte, 75006 Paris, France; <nils-robin@orange.fr>). Il est recommandé aux adhérents de la Zone Euro, hors France, de procéder par virement bancaire SEPA (gratuit), l'encaissement des chèques émis hors de France étant très coûteux. Un RIB leur sera adressé sur demande. Également, les cotisations peuvent être payées en ligne sur notre site internet <<http://malimbus.free.fr/>>. Les Membres Ordinaires reçoivent *Malimbus* par courrier ordinaire et les Sociétés par courrier aérien, gratuitement. Un supplément est exigé des Membres Ordinaires pour le courrier aérien (demander au Trésorier le tarif).

Numéros anciens: Aucun exemplaire original n'est disponible à l'exception de quelques numéros récents. Cependant, un pdf de l'un des numéros (numéro complet) de *Malimbus* ou du *Bull. Nigerian Orn. Soc.* peut être fourni au prix de 50 % du coût de l'abonnement de l'année en cours. Merci d'adresser vos requêtes au Trésorier.

Bourses de Recherches de la S.O.O.A.: Les conditions à remplir pour les candidatures se trouvent dans *Malimbus* 25: 72-75 et sur le site web, ou peuvent être obtenues auprès du Secrétaire du Conseil (adresse ci-dessus).



Editorial

Our 50th Anniversary

The beginnings of W.A.O.S. came when a group of enthusiasts created the Nigerian Ornithologists' Society, in 1964, so this year marks our 50th birthday. In the early days, almost all of the contributions to the new journal, the *Bulletin of the Nigerian Ornithologists' Society*, came from expatriates. It is thus gratifying to note several features of the Society and its journal after 50 years. Above all, we still exist, despite a couple of periods when it looked as if *Malimbus* might die; but the Society maintains its relevance, and its journal has become a prime repository for research on the birds of more than one third of the continent. Secondly, the articles published in the journal have come to include an increasing proportion written by West African authors. This is a measure of success in one of the Society's objectives: to encourage and promote research by West Africans in West Africa.

Several features of the present issue in particular are perfectly appropriate to mark our golden jubilee. The first is that in this issue all the full-length articles are written by West African authors (the first time this has occurred) and, given that the anniversary is of 50 years since our founding as the *Nigerian Ornithologists' Society*, it is doubly appropriate that two of these articles should come from Nigeria. Further, one of them examines the changes in the avifauna of the Ibadan area in the 50 years since our foundation, which happened to coincide with the publication of a key work on the district by John Elgood (one of our founders) and Frank Sibley. The other re-examines aspects of the avifauna of the Yankari Game Reserve, which N.O.S. and W.A.O.S. members have studied since our beginnings. The third article, from francophone West Africa, demonstrates that our move to a bilingual journal, when the Society's geographical coverage expanded to the whole of West Africa, is still valued. And finally, all this is pure coincidence: for anyone inclined to believe in augury, a set of auspicious signs for the future from West Africa's birds.

The continuation of contributions from Nigeria is in part due to the stimulus that is still provided by old-established scholarly institutions there, including the university and the International Institute of Tropical Agriculture at Ibadan, along with the relatively young A.P. Leventis Ornithological Research Institute. We can also recognize with satisfaction that N.O.S. and W.A.O.S. have contributed to this stimulus.

So, despite some worrying periods for the Society and its journal in the past (and still, with submissions this year very low), we have good reason to expect ornithology in Nigeria and West Africa to continue, with its practitioners providing valuable new knowledge and contributing to the conservation of West Africa's birds. Surely enough reason to celebrate.

Alan Tye

50ème anniversaire !

Les débuts de la S.O.O.A. remontent à la création en 1964 de la Nigerian Ornithologists' Society par un groupe d'enthousiastes, et donc cette année marque notre 50ème anniversaire. Au début, la presque totalité des contributions au nouveau journal, le *Bulletin of the Nigerian Ornithologists' Society*, provenait d'expatriés. Il est ainsi gratifiant de relever, après 50 ans, plusieurs faits saillants concernant notre Société et son journal. Avant tout, nous existons encore, malgré deux périodes au cours desquelles *Malimbus* put sembler sur le point de disparaître; mais la société conserve sa raison d'être et son journal est devenu l'une des principales revues scientifiques sur l'avifaune de plus d'un tiers du continent africain. En second lieu, une proportion croissante des articles publiés dans le journal ont été écrits par des auteurs d'Afrique de l'Ouest. C'est l'une des mesures de la réussite de la Société dans l'atteinte de l'un de ses objectifs: encourager et promouvoir la recherche en Afrique de l'Ouest par des Africains de l'Ouest.

Plusieurs aspects de ce numéro sont particulièrement bienvenus pour marquer notre jubilé. Le premier est que dans ce numéro, pour la première fois, tous les articles longs ont été écrits par des auteurs Ouest Africains. D'autre part, s'agissant du 50ème anniversaire depuis notre création en tant que *Nigerian Ornithologists' Society*, il est d'autant plus opportun que deux de ces articles proviennent du Nigeria. De plus, l'un d'eux examine ce qui a changé dans l'avifaune d'Ibadan depuis 50 ans, alors précisément que notre création avait coïncidé avec la publication d'un travail clé sur le district par John Elgood, l'un de nos fondateurs, et Frank Sibley. L'autre article réexamine des aspects de l'avifaune de la Réserve de faune de Yankari, que des membres de N.O.S. et de la S.O.O.A. avaient étudiés depuis nos débuts. Le troisième article, d'Afrique de l'Ouest francophone, démontre que notre évolution vers un journal bilingue, alors que la couverture géographique de la Société s'était étendue à toute l'Afrique de l'Ouest, conserve toute sa valeur. Et enfin, tout ceci est une pure coïncidence: pour tous ceux enclins à croire aux présages, voilà peut-être un ensemble de signes prometteurs quant à l'avenir des oiseaux d'Afrique de l'Ouest.

Le fait que nous continuions à recevoir des contributions du Nigeria est en partie dû à la stimulation encore exercée par des institutions d'installation ancienne, dont l'Université et l'Institut International d'Agriculture Tropicale d'Ibadan, ainsi que l'Institut de Recherche Ornithologique A.P. Leventis. Nous pouvons également considérer avec satisfaction que la N.O.S. et la S.O.O.A. ont contribué à cette stimulation.

Ainsi, bien que la Société et son journal aient connu des périodes inquiétantes dans le passé (et encore aujourd'hui, avec des soumissions très peu nombreuses cette année), nous avons de bonnes raisons d'avoir confiance dans l'avenir de l'ornithologie au Nigeria et en Afrique de l'Ouest, avec des praticiens produisant de précieuses nouvelles connaissances et contribuant à la conservation des oiseaux en Afrique de l'Ouest. À coup sûr, des raisons suffisantes pour célébrer cet anniversaire.

Alan Tye (traduit: Nils Robin)

Raptors in Yankari Game Reserve and surrounding unprotected area, Nigeria

by J.D. ONOJA^{1,2}, T. TENDE^{1,3}, T.C. OMOTORIOGUN^{1,4}, U. OTTOSSON¹,
S.A. MANU^{1,2} & G.S. MWANSAT²

¹A.P. Leventis Ornithological Research Institute, PO Box 13404, Jos Plateau State, Nigeria

²Dept of Zoology, University of Jos, PMB 2084, Jos, Nigeria <oj_daniels@yahoo.com>

³Dept of Animal Ecology, Lund University, Sweden

⁴National Centre for Biosystematics, Natural History Museum, Univ. of Oslo, Norway

Received 15 May 2012; revised 27 February 2014.

Summary

A recent survey to estimate the population of raptors in and around Yankari Game Reserve revealed that the reserve still holds a number of raptor species also found in other reserves in West Africa. In total, 886 raptors of 37 species were recorded in the gallery forest and savanna habitats of the reserve, while 155 individuals in 18 species were encountered outside of the reserve. The number of raptors per km of transect in the unprotected area was lower than in the reserve, despite transect speed being slower and raptor detectability probably higher in the unprotected area. Most of the species encountered during surveys were medium-sized raptors. Juvenile raptors were seen only within the reserve, and adults of large species were rare outside it. Vultures seem to have declined since 2008, with two species having apparently been lost from the area.

Résumé

Les rapaces dans la Réserve Naturelle de Yankari et la zone adjacente non protégée, Nigeria. Une étude récente visant à estimer la population de rapaces dans et autour de la Réserve Naturelle de Yankari a révélé que la réserve contient encore nombre d'espèces de rapaces également présentes dans d'autres réserves d'Afrique de l'Ouest. Au total, 886 rapaces de 37 espèces ont été observés dans les habitats de forêt galerie et de savane, tandis que 155 individus de 14 espèces ont été rencontrés hors de la réserve. Le nombre de rapaces par km de transect dans la zone non protégée était inférieur à celui de la réserve, bien que la vitesse de parcours des transects ait été plus lente et la possibilité de découverte de rapaces probablement plus forte dans la zone non protégée. La plupart des espèces rencontrées au cours des enquêtes

ont été des rapaces de taille moyenne. Des rapaces juvéniles n'ont été vus qu'à l'intérieur de la réserve et les adultes d'espèces de grande taille étaient rares à l'extérieur. Les vautours paraissent être moins nombreux depuis 2008, avec deux espèces ayant apparemment disparu de la zone.

Introduction

Diurnal birds of prey are good indicators of changes in ecosystems and of impacts of human activity, owing to their sensitivity to environmental contamination, persecution and disturbance (Newton 1979). Most species of raptor are conspicuous and they feed on a broad array of invertebrates and vertebrates across all natural and artificial habitats (Thiollay 2006). Today, the dramatic increase in human pressure, deforestation and general decrease in tree cover, overgrazing and erosion, pesticide use, over-hunting and fishing, and sometimes direct persecution of predators, all impact negatively on raptor population (Thiollay 2006). To counter widespread population declines, many attempts have been made in recent years to increase raptor numbers, either by the management of the birds themselves or of their habitat and food sources (Newton 1979). The positive effects on these birds are revealed in nature reserves, where threats to them are minimized. For example in West Africa an estimate of the mean abundance index of raptors between protected and unprotected areas has shown a 30 % decline of raptors in protected areas compared with a 67 % decline in unprotected areas (Thiollay 2006, 2007b). In the savannas of West Africa, 40 breeding raptor species and 19 Palearctic migrant species have been recorded (Borrow & Demey 2001, Thiollay 2007a). Brown (1970) observed that little is known about the actual numbers of the various species, and without knowing this it is difficult to determine the real effect of conservation effort.

A recent survey in Yankari Game Reserve, Bauchi State, Nigeria, revealed a decline in the population of vultures (Tende & Ottosson 2008). However, data on the population status of other raptor species is lacking. The aim of these surveys was therefore to estimate the abundance, diversity and distribution of raptors in Yankari Game Reserve and surrounding unprotected areas, in a bid to establish a baseline for conservation for this group of birds.

Methods

Study site

The survey was carried out in and around the Yankari Game Reserve (9°50'N, 10°30'E), Bauchi State in the east-central part of Nigeria. The reserve has a total area of 2244 km² and is bisected by the River Gaji. It lies within the Sudan Savanna Zone (Geerling 1973) and receives an average rainfall of *c.* 1000 mm per year, between

April and October (Crick & Marshall 1981). The vegetation includes swampy flood plain bordered by patches of forest, gallery forest, riparian forest and woodland savanna (Crick & Marshall 1981, Ezealor 2002). In seasonally flooded “fadamas”, *Ficus* spp. and *Mitragyna* sp. are the dominant trees, while tangles of *Mimosa pigra* dominate the shrub stratum. The reserve was established as a Game Preservation Unit (GPU) of the Northern Region of Nigeria in January 1956 (Sikes 1964), and from its inception has been traversed by roads, by which border zone communities reach and utilize renewable natural resources within the reserve. Presently, twelve communities around the reserve have roads leading to its centre at Wikki Camp (9°45'N, 10°30'E). About 337 species of birds have been recorded (Ezealor 2002), of which 130 are resident, 50 are Palearctic migrants and the rest are intra-African migrants.

The unprotected area outside the reserve is mainly farmland with scattered trees and hedges of *Guerra senegalensis*. Crops grown in these farmlands were mostly groundnut *Arachis hypogaea*, beans *Phaseolus vulgaris*, guinea corn *Sorghum bicolor*, millet *Pennisetum glaucum* and rice *Oryza sativa*.

Survey Design

The survey within the reserve was carried out for eight weeks from 21 April to 28 June 2008. This period coincided with the beginning of the rainy season, with insignificant rainfall during the period of the survey. Twenty line-transects (Bibby *et al.* 2000) of an average length of 10 km were used to count raptors. A total distance of 1636 km was covered during this transect survey. Transects were along game viewing and anti-poaching patrol tracks, and chosen to cover both savanna and gallery forest habitats. The coordinates of the start and end of each transect were recorded. Surveys were carried out in the morning (6h30–13h00) and evening (15h00–18h00). Counts were made by two non-driving observers from a car moving at an average speed between 12 and 25 km/h (Fuller & Mosher 1981). All raptors flying, perched and heard were identified, counted and their locations recorded using a Garmin 760CSx global positioning system (GPS) receiver. For birds seen perched, their perpendicular distances from the transect were recorded using a Canon Laser Range Finder.

In addition, observations were made from five different vantage points with good views of the surroundings (Fig. 1). Each point was visited at least three times and one hour was spent at each point, between 12h00 and 13h00, *i.e.* at a time of day when raptors make use of thermals to soar. Using binoculars and a telescope, the surroundings were scanned for any raptors seen perched or flying. Efforts were made to avoid double counts of birds observed by noting the direction in which the last counted individual disappeared from sight and the flock size.

The unprotected area outside the reserve was surveyed between 23 May and 23 July 2009. A total of 33 transects, each measuring 10 km in length, was laid in the area surrounding the reserve. Counts were made by two non-driving observers from a car moving at an average speed of 10.5 km/h (slower than in the reserve because of the bumpy nature of the roads). All raptors seen perched or flying or heard were




Figure 1. LandSat photograph of Yankari Game Reserve and surrounding unprotected area, showing observation vantage points.

identified and their location recorded. Surveys were carried out between 10h00 and 14h00, and all transects were visited once. The survey covered a total distance of 330 km. Five vantage points outside the reserve were used for observing raptors soaring or perched (Fig. 1).

Data Analysis

The surveys were designed to provide baseline data for each of the two areas and not primarily for comparison between them. Detectability, timing of survey periods and speed of the vehicle were all different in the two habitats. Detectability was probably higher in the unprotected area than in the reserve because of the more open nature of the unprotected area, and transect speed was slower there. This must be borne in mind when using these data for comparison between the two habitats.

Abundance on transects, of all raptors combined and of each raptor species, is expressed as encounter rate, in terms of the number of individuals encountered per 100 km of transect travelled and per h of transect time. Encounter rate during vantage point observations is expressed as number of raptors seen per h of observation. The encounter rates on transects were calculated as total number of individuals encountered over the total distance covered per month during the period of survey.

Results

In the reserve, 886 individual raptors of 37 species were recorded during the transects, combining forest and savanna (Table 1), of which 49 individuals (5.5 %) were classed as juveniles and sub-adults. In April, 170 raptors were seen over a total distance of 301.1 km covered (56 birds per 100 km), May had 418 raptors over 789.1 km (53 per 100 km), while June had 298 raptors over 545.3 km (55 per 100 km): there was no significant difference in encounter rate on transects between months ($\chi^2 = 0.0854$, $df = 2$, $P = 0.958$). The overall raptor encounter rates during transects in the reserve were 54 birds per 100 km (total 1635.6 km) or 5.5 birds per h (total 160.35 h), with higher overall encounter rate in the forest than in the savanna (Table 1). In the savanna, the Grasshopper Buzzard was the species with the highest encounter rate, with other common species including Dark Chanting-Goshawk, Grey Kestrel and Bateleur, while the Palmnut Vulture had the highest encounter rate for the gallery forest with other commonly encountered species there including Bateleur, African Fish-Eagle, Black Kite, Lizard Buzzard, Grey Kestrel and African Hawk-Eagle (Table 1). Encounter rates at vantage point observations in the reserve were higher than the per h results from transects, but because less time was spent on the former (total 17 h of observation), the results for each species are less informative and are not presented here; 126 raptors were counted during vantage point observations in the reserve, giving an overall encounter rate of 7.4 birds/h. During the ten weeks of fieldwork, the Secretary Bird *Sagittarius ser-pentarius* was sighted only once, in the reserve but outside the survey period on 21 April.

For the unprotected area, 155 individual raptors, of 18 species, were encountered during the transect survey. In May, 36 raptors were seen and a total distance of 70 km was covered (51 birds per 100 km), June had 105 raptors over 170 km (62 per 100 km), while July had 14 raptors over 90 km (16 per 100 km). There was a significant difference between months in the encounter rates of raptors during the transect survey ($\chi^2 = 26.84$, $df = 2$, $P < 0.0001$), largely because Grasshopper Buzzard was common in May and June but not recorded in July. The overall raptor encounter rate during transects in the unprotected area was 47 birds per 100 km (c. 330 km) and 4.5 birds per h (34.5 h of survey). The species with the highest encounter rate was Black Kite, while Grasshopper Buzzard and Fox Kestrel were also common (Table 1). African Swallow-tailed Kite, Bateleur, Brown Snake Eagle, Eurasian Marsh Harrier, Lanner and Lizard Buzzard were encountered only once each, the Eurasian Marsh Harrier in a rice field on 23 May. During vantage point observations outside the reserve, the encounter rate was 3.5 birds per h (26 birds over 7.5 h of observation).

Discussion

This survey shows that the Yankari Game Reserve still retains a considerable number of raptor species that normally occur in other nature reserves in West Africa (*cf.* Thiollay

Table 1. Transect counts of raptor species in savanna and gallery forest in the Yankari Game Reserve, and in the surrounding unprotected area.

| | Savanna | | Gallery | | Unprotected | |
|---|-------------|-------------|-------------|-------------|-------------|-------------|
| | total count | n/100 km | total count | n/100 km | total count | n/100 km |
| <i>Pandion haliaetus</i> Osprey | - | - | 2 | 0.34 | - | - |
| <i>Pernis apivorus</i> Western Honey-Buzzard | 2 | 0.19 | 2 | 0.34 | - | - |
| <i>Macheiramphus alcinus</i> Bat Hawk | - | - | 1 | 0.17 | - | - |
| <i>Elanus caeruleus</i> Black-shouldered Kite | 18 | 1.73 | 4 | 0.67 | 10 | 3.03 |
| <i>Chelictinia riocourii</i> Swallow-Tailed Kite | - | - | - | - | 1 | 0.30 |
| <i>Milvus migrans</i> Black Kite | 11 | 1.06 | 34 | 5.71 | 41 | 12.42 |
| <i>M. parasitus</i> Yellow-billed Kite | 2 | 0.19 | 2 | 0.34 | - | - |
| <i>Haliaeetus vocifer</i> African Fish-Eagle | 5 | 0.48 | 48 | 8.07 | - | - |
| <i>Gypohierax angolensis</i> Palmnut Vulture | 8 | 0.77 | 61 | 10.25 | - | - |
| <i>Necrosyrtes monachus</i> Hooded Vulture | - | - | 5 | 0.84 | - | - |
| <i>Gyps africanus</i> African White-backed Vulture | 1 | 0.10 | 13 | 2.35 | - | - |
| <i>Trigonoceps occipitalis</i> White-headed Vulture | 8 | 0.77 | 4 | 0.67 | - | - |
| <i>Circaetus gallicus</i> Short-toed Snake-eagle | - | - | 1 | 0.17 | - | - |
| <i>C. cinereus</i> Brown Snake-eagle | 7 | 0.67 | 8 | 1.34 | 1 | 0.30 |
| <i>C. cinerascens</i> Western Banded Snake-Eagle | 1 | 0.10 | 1 | 0.17 | - | - |
| <i>Terathopius ecaudatus</i> Bateleur | 36 | 3.46 | 49 | 8.23 | 1 | 0.30 |
| <i>Polyboroides typus</i> African Harrier-Hawk | 15 | 1.44 | 14 | 2.35 | - | - |
| <i>Circus macrourus</i> Pallid Harrier | 3 | 0.29 | - | - | - | - |
| <i>C. ranivorus</i> African Marsh Harrier | 1 | 0.10 | - | - | - | - |
| <i>C. aeruginosus</i> Eurasian Marsh Harrier | - | - | - | - | 1 | 0.30 |
| <i>Melierax metabates</i> Dark Chanting-Goshawk | 56 | 5.38 | 3 | 0.50 | 11 | 3.33 |
| <i>Accipiter badius</i> Shikra | 13 | 1.25 | 16 | 2.69 | 6 | 1.82 |
| <i>A. ovampensis</i> Ovambo Sparrowhawk | - | - | 2 | 0.34 | - | - |
| <i>Butastur rufipennis</i> Grasshopper Buzzard | 183 | 17.59 | 12 | 2.02 | 36 | 10.91 |
| <i>Kaupifalco monogrammicus</i> Lizzard Buzzard | 19 | 1.83 | 29 | 4.87 | 1 | 0.30 |
| <i>Buteo auguralis</i> Red-necked Buzzard | 7 | 0.67 | 1 | 0.17 | 11 | 3.33 |
| <i>Aquila rapax</i> Tawny Eagle | 3 | 0.29 | 4 | 0.67 | - | - |
| <i>A. wahlbergi</i> Wahlberg's Eagle | 1 | 0.10 | 3 | 0.50 | 2 | 0.61 |
| <i>Hieraaetus spilogaster</i> African Hawk-Eagle | 15 | 1.44 | 22 | 3.70 | 2 | 0.61 |
| <i>Lophaaetus occipitalis</i> Long-crested Eagle | 1 | 0.10 | 3 | 0.50 | 4 | 1.21 |
| <i>Polemaetus bellicosus</i> Martial Eagle | 2 | 0.19 | 4 | 0.67 | - | - |
| <i>Falco alopex</i> Fox Kestrel | 3 | 0.29 | -- | -- | 18 | 5.45 |
| <i>F. ardosiaceus</i> Grey Kestrel | 40 | 3.84 | 25 | 4.20 | 5 | 1.52 |
| <i>F. chicquera</i> Red-necked Falcon | 5 | 0.48 | 7 | 1.18 | 3 | 0.91 |
| <i>F. vespertinus</i> Red-footed Falcon | 11 | 1.06 | - | - | - | - |
| <i>F. cuvierii</i> African Hobby | 4 | 0.38 | 7 | 1.18 | - | - |
| <i>F. biarmicus</i> Lanner Falcon | 4 | 0.38 | 2 | 0.34 | 1 | 0.30 |
| <i>Glauclidium perlatum</i> Pearl-spotted Owlet | - | - | 2 | 0.34 | - | - |
| Totals | 485 | 46.6 | 391 | 65.9 | 155 | 47.0 |

2006). However, only three species of vulture were seen during the survey, in contrast to the five species recorded by Tende & Ottosson (2008), who also found Lappet-faced Vulture *Torgos tracheliotos* and European Griffon *Gyps fulvus*. The population of vultures in the reserve has declined markedly as compared with earlier surveys (Dyer & Gartshore 1975, Crick & Marshall 1981). Similarly, in the Sudan Savanna zone of West Africa, Thiollay (2006) reported a decline of 97 % of the African White-backed Vulture in 30 years. Possible reasons for such declines may be poisoning and persecution, and Thiollay (2007a) suggested that habitat degradation was a major factor. Protected areas remain a haven for raptors and should be maintained as such. However, the decline of vulture populations observed in the reserve may be due to poisoning outside the protected area (Tende & Ottosson 2008).

Other large raptor species such as Martial Eagle, Long-crested Eagle, Tawny Eagle, African Hawk Eagle and Wahlberg's Eagle were recorded in the reserve, including juveniles of Bateleur, Palmnut Vulture, African Hawk-eagle and Martial Eagle, suggesting that it may be an important breeding ground for raptor species. During both vantage point observations and road transects outside the reserve, only adult birds were seen. It may be that the species recorded outside the reserve mostly breed within it and only use the unprotected area for hunting.

The single sighting of Secretary Bird contrasts with the situation in the 1970s, when Demeter (1977) suggested that the bird occurred regularly in the reserve in low numbers. However, this species has declined so much that it is not seen in Nigeria now outside protected areas and only rarely within them.

The species encountered most often outside the reserve were of medium size: Black Kite, Fox Kestrel, Grasshopper Buzzard (in May–June), Dark Chanting-Goshawk and Red-necked Buzzard. The Fox Kestrel was mostly found in an area west of the reserve, probably due to the rocky nature of the habitat there (*cf.* Brown *et al.* 1997). Among the larger raptors recorded in the unprotected area were Bateleur (seen once, soaring) and Long-crested Eagle (seen twice outside and twice soaring over the protected area). Larger raptors may not find the unprotected areas conducive because of inadequate food supply and nest sites due to the degraded nature of the habitat. African Swallow-tailed Kite was recorded once outside the reserve; this species has never been recorded within the reserve, either during this survey or previously. The largest raptor encountered regularly in open plains and farmlands was the Red-necked Buzzard. No vultures were encountered in the unprotected area, perhaps because of the severe persecution that vultures and other raptors have suffered (*e.g.* Rondeau & Thiollay 2004, Thiollay 2006). When seen, vultures were either soaring very high or within the reserve.

There was little difference in the number of birds recorded between months probably because most of the species recorded are residents or have resident populations. In the case of Grasshopper Buzzard, not recorded in July during the survey outside the reserve, nor within it that month (casual observations), it may have migrated northwards to breed then (Brown *et al.* 1997).

The average number of raptors observed in the unprotected area was 0.48 birds/km, lower than in the protected area (0.75), despite the transect speed being slower and raptor detectability probably higher outside the reserve. It is also lower than in other studies (1.63, Thiollay 2006; 0.58, Tende & Ottosson 2008). The lower rate outside the reserve could be partly due to the surveys there having been carried out later than those in the reserve, so that Palaearctic migrants such as Honey-Buzzard, harriers and Red-footed Falcon would have departed by the later part of the unprotected area survey. However, higher encounter rates were found outside the reserve for the migrant Black Kite and, overall, the proportion of species encountered that were Palaearctic migrants was small, so that this factor could not account for the generally lower encounter rate in the unprotected area. Rather, other factors such as pesticide use on farmlands may have reduced the number of insects and rodents in unprotected areas, making them less suitable for raptors as compared with protected areas. Overgrazing and cutting of trees and other vegetation for farming have also resulted in low quality habitat in the unprotected area, and the area is highly disturbed by people. In unprotected areas there may also be direct persecution of raptors. People sometimes kill them for meat or to prevent them from preying on their domestic chickens (Tende & Ottosson 2008). These reasons may be the cause of the decline of raptors, especially the large ones, while the medium-sized species that can survive on small prey like insects, scorpions and rodents are relatively less affected.

Although the reserve's population of vultures and other raptor species has declined, it still has the potential to sustain the populations of raptors left, its habitat heterogeneity making it a good haven for many species of raptors. The reserve also has great potential as an ecotourism destination and a base for biological research. Therefore, it is recommended that efforts should be made to enlighten the surrounding communities on the need to stop the persecution of raptors. A raptor monitoring programme also needs to be put in place to detect changes both within and outside the reserve.

Acknowledgments

This work for the protected area was generously funded by the Leventis Foundation. The authors are deeply grateful to the British Ornithologists' Union for providing additional funds and to the Swedish Ornithological Society for support. We appreciate Haladu Idi's support in the field. This is contribution no. 74 from the A.P. Leventis Ornithological Research Institute, to which we are grateful for logistical support.

References

- BIBBY, C.J., BURGESS, N.D., HILL, D.A. & MUSTOE, S.H. (2000) *Bird Census Techniques*. Academic Press, London.

- BORROW, N. & DEMEY, R. (2001) *Birds of Western Africa*: Christopher Helm, London.
- BROWN, L., URBAN, E.K. & NEWMAN, K. (1997) *The Birds of Africa*, vol. 2. Academic Press, London.
- BROWN, L. (1970) *African Birds of Prey*. Collins, London.
- CRICK, H.Q.P. & MARSHALL, P.J. (1981). The birds of Yankari Game Reserve, Nigeria: their abundance and seasonal occurrence. *Malimbus* 3: 103–113.
- DEMETER, A.A. (1977) Additions to local avifaunas: Yankari. *Bull. Niger. Orn. Soc.* 13: 81.
- DYER, M. & GARTSHORE, M.E. (1975) Birds of Yankari Game Reserve, Nigeria. *Bull. Niger. Orn. Soc.* 11: 77–84.
- EZEALOR A.U. (ed.) (2002) *Critical Sites for Biodiversity Conservation in Nigeria*. Nigerian Conservation Foundation, Lagos.
- FULLER, M.R., & MOSHER, J.A. (1981) Methods of detecting and counting raptors: a review. *Stud. avian Biol.* 6: 235–246.
- GEERLING, C. (1973) *The Vegetation of Yankari Game Reserve, its Utilization and Condition*. Bulletin 3, Dept of Forestry, University of Ibadan, Ibadan.
- NEWTON, I. (1979) *Population Ecology of Raptors*. Poyser, Berkhamsted.
- RONDEAU, G. & THIOLLAY, J.M. (2004) West African vulture decline. *Vulture News* 51: 13–33.
- SIKES, S.K. (1964) A game survey of Yankari Game Reserve, Northern Nigeria. *Niger. Field* 36: 171–180.
- TENDE, T. & OTTOSSON, U. (2008) The current status of vultures in Yankari. *Vulture News* 59: 7–12.
- THIOLLAY, J.M. (2006) The decline of raptors in West Africa: long-term assessment and the role of protected areas. *Ibis* 148: 240–254.
- THIOLLAY, J.M. (2007a) Raptor population decline in West Africa. *Ostrich* 78: 405–413.
- THIOLLAY, J.M. (2007b) Raptor declines in West Africa: comparisons between protected, buffer and cultivated areas. *Oryx* 41: 1–8.

Birds of the International Institute of Tropical Agriculture campus, a stronghold of avian diversity in the changing Ibadan area (Nigeria) over the last 50 years

by Taiye A. ADEYANJU^{1,2,3}, Ulf OTTOSSON¹, Temidayo E. ADEYANJU¹,
Taiwo C. OMOTORI OGUN^{1,4}, Philip HALL¹, Shiiwua A. MANU¹, Tunrayo ALABI⁵,
Gbolagade A. LAMEED² & Deni BOWN⁵

¹A.P. Leventis Ornithological Research Institute,
Laminga, Jos Plateau State, P.O. Box 1304, Nigeria

²Department of Wildlife and Ecotourism Management, Univ. of Ibadan, Nigeria

³Correspondence: <taiyelongifolia@yahoo.com>

⁴National Centre for Biosystematics, Natural History Museum, Univ. of Oslo, Norway

⁵International Institute of Tropical Agriculture, Ibadan, Nigeria

Received 20 August 2012; revised 14 May 2014.

Summary

Data from a survey carried out between 2009 and 2013 of the bird species in the International Institute of Tropical Agriculture (IITA) campus, Ibadan, Nigeria, are compared with records from the area around Ibadan (including IITA) for the last 50 years. The IITA campus is an Important Bird Area (IBA) and includes a secondary dry semi-deciduous forest reserve of c. 360 ha, which is fenced and protected. The forested area is now being extended and enhanced by planting indigenous trees on degraded farm plots, and by enrichment planting in degraded areas of forest. In total, 398 bird species from at least 71 families are now recorded for the Ibadan area, of which 322 species have been recorded since 2002. The IBA holds at least 269 of these species (68 %) in 64 families, while the forest reserve holds c. 137 species. Seventy-five species were mis-netted in the IBA in our study. Twenty-five species plus 13 vagrants are new to the IBA and the Ibadan area in general, having not been detected prior to 2002. However, a minimum 68 species plus an additional 62 vagrant species reported in the Ibadan area by earlier studies have not been detected recently. We report an additional 29 biome-restricted species present in the Ibadan area (74 had been reported previously), 17 of which occur in the IBA. Diversity of some groups of large birds (e.g. Anatidae) has declined whereas many forest edge or generalist species (e.g. Double-spurred Francolin *Francolinus bicalcaratus*) have increased in abundance and range. Forest specialists, including many Pycnonotidae and Bucerotidae species, appear to have declined. The IITA campus, with its forest reserve, lakes and farm plots, is now an “island” IBA

of great avifaunal diversity surrounded by a highly modified anthropogenic landscape unwelcoming for many of the birds that formerly inhabited the area.

Résumé

Les oiseaux du campus de l'Institut International d'Agriculture Tropicale, un bastion de la diversité de l'avifaune dans la région d'Ibadan (Nigeria), en évolution rapide au cours des 50 dernières années. Les données d'une étude réalisée entre 2009 et 2013 sur les espèces d'oiseaux dans le campus de l'Institut International d'Agriculture Tropicale (IIAT), Ibadan, Nigeria, sont comparées avec des observations dans les alentours d'Ibadan (incluant l'IIAT) pendant les 50 dernières années. Le campus de l'IIAT est une Zone Importante pour la Conservation des Oiseaux (ZICO) et inclut une réserve de forêt secondaire sèche semi-décidue de c. 360 ha, clôturée et protégée. Cette zone de forêt est en cours d'extension et d'amélioration par la plantation d'espèces d'arbres indigènes sur des terrains agricoles dégradés et par des plantations visant à enrichir des zones dégradées de la forêt. Au total, 398 espèces d'oiseaux appartenant à au moins 71 familles sont aujourd'hui recensées pour la zone d'Ibadan, dont 322 espèces ont été observées depuis 2002. La ZICO héberge au moins 269 de ces espèces (68 %) en 64 familles, cependant que la réserve forestière en héberge c. 137 espèces. Soixante-quinze espèces ont été capturées au filet dans la ZICO au cours de notre étude. Vingt-cinq espèces plus 13 occasionnelles sont nouvelles pour la ZICO et la zone d'Ibadan en général, n'ayant pas été observées avant 2002. Cependant, au moins 68 espèces plus 62 espèces occasionnelles observées dans la zone d'Ibadan lors d'études antérieures n'ont pas été observées récemment. Nous mentionnons 29 espèces restreintes à un seul biome qui sont nouvelles dans la zone d'Ibadan (74 avaient été précédemment mentionnées), parmi lesquelles 17 sont présentes dans la ZICO. La diversité de quelques groupes de grands oiseaux (p. ex. Anatidae) a décliné alors que de nombreuses espèces péri-forestières ou généralistes (p.ex. Francolin à double éperon *Francolinus bicalcaratus*) a augmenté en abondance et zones de présence. Les espèces forestières, dont de nombreux Pycnonotidae et Bucerotidae, apparaissent en déclin. Le campus de l'IIAT, avec sa réserve forestière, des lacs et des terrains agricoles, est maintenant une "île" ZICO d'une grande diversité avifaunistique au milieu d'un paysage complètement modifié par l'homme et devenu inhospitalier pour de nombreux oiseaux autrefois présents dans la zone.

Introduction

The campus of the International Institute of Tropical Agriculture (IITA) at Ibadan (7°30'N, 3°55'E) occupies c.1000 ha (Fig. 1), including a 360 ha relict of secondary dry




Figure 1. The IITA campus.

semi-deciduous rainforest, the “forest reserve”. The rest of the campus, apart from the residential buildings, workshops and offices, consists of lakes, rice paddies, farm plots, marshes and bushes, which provide additional habitats for wildlife. The forest area on the campus is now being extended by reforestation and therefore the term “forest land use” on the map (Fig. 1) includes areas preserved as forest (since 1965) and areas recently reforested (during the last four years). The bird species at IITA include aquatic, savanna and forest species, and the campus has been recognised as an Important Bird Area (IBA: Ezealor 2001, A. Ezealor pers. comm.). This protected ecosystem on the IITA campus has encouraged visitors, research and collaboration; nonetheless it has also encouraged poachers, though hunting is prohibited on the site. However there is no detailed, published and up to date checklist of the avifauna of either the IITA campus as a whole (the IBA) or the forest reserve.

The Ibadan area was described in detail by Elgood & Sibley (1964). It has since grown into a large city where demand for fuel wood, timber for furniture and building, farmland and other uses has degraded the natural habitat of the area. Most areas mentioned by Elgood & Sibley (1964) and others (see Table 1 header) as holding high bird

diversity are now a shadow of their former selves. However in the midst of this changing environment, the IITA campus retains a rich diversity of resident and migratory birds.

Ibadan is located in an area divided between derived savanna and forest ecosystems. The IITA campus possesses areas of both these ecosystems, in the form of degraded farmland and secondary forest respectively. Early studies on the avifauna of the Ibadan area are numerous, though many were carried out prior to the establishment of the IITA headquarters in 1965, and the consequent protection of the site. One of the most complete early studies, which covered the general area in which IITA is now situated, was that on bird species distribution in Ibadan and southwest Nigeria by Elgood & Sibley (1964). This summarised earlier records for the Ibadan area, defined as within a 10-mile (16 km) radius of Mapo Hall at the centre of the city, and with an emphasis on birds found within and adjacent to the extensive grounds of the University of Ibadan (UI). Elgood & Sibley (1964) reported 266 bird species in total, and showed that the terrestrial element in the avifauna was evenly balanced between forest and savanna forms. The IITA and UI are < 7 km apart, separated by villages, though tracts of semi-natural vegetation are present. Nonetheless, some birds are capable of flying between the two sites daily as observed for Cattle Egrets and White-faced Whistling Ducks, which roost at UI and IITA respectively (pers. obs.). Also, the checklist of the birds of Nigeria (Elgood *et al.* 1994) gives numerous early records for Ibadan.

Since the establishment of the IITA campus, there have been only a few published records from within it. The only study specifically on IITA birds was by Ezealor (2001) for IBA designation (74 spp.). A single record for IITA was given by Ash (1990), 17 records by Elgood *et al.* (1994), seven more by Demey *et al.* (2003), and 29 others from a study of the effects of forest fragmentation on the endangered Ibadan Malimbe *M. ibadanensis* (Manu *et al.* 2005). Among earlier and more recent unpublished records within IITA are PH's from casual walks through the area during the last 25 years, and others listed in the header to Table 1. However, there remained a need for a thorough survey of the forest reserve and its environs within the IITA campus, to bring up to date and consolidate our knowledge of the bird species composition of the site. Our study therefore aims to provide an updated checklist of the avifauna at IITA. We also compare recent records from the IITA campus with earlier published and unpublished reports from within it, in order to investigate changes in the avifauna of the campus during the past 25 years. We also compare these records with earlier and more recent records from the surrounding Ibadan area, particularly UI, Eleyele, Ibadan Golf Club and Moor Plantation in order to determine how much of the diversity previously known to occur in Ibadan is being maintained by the protection of the IITA campus since 1965.

Methods

Each month from 2009 to 2012 counts were made along nine transects, each of 1 km, of which three were placed along pre-cut walking trails in each of farmland, forest and

along the shore of the lake (forest and lake-shore transects shown on Fig. 2). In addition, counts were made from vantage points at rice paddies and lakes within IITA. Visits were made once each month of the year, all in the mornings between 6h00 and 12h00 and evenings between 15h00 and 19h00). On each visit, two or three transects were surveyed by TAA, walking at an average speed of *c.* 1 km/h (open habitats) or *c.* 0.5–0.7 km/h (forest interior, due to lower detection rates and need to spend more time on bird calls). Birds seen outside the transect times are also included. General abundance of species sighted during our study is classed as: Very Abundant (VA) >100 may be seen or heard in suitable habitat per day; Abundant (A) 11–100 may be seen or heard in suitable habitat per day; Common (C) 1–10 may be seen or heard in suitable habitat per day; Frequent (F) often seen but not every day; Uncommon (U) several records per year; Rare (R) one record per several years (resident species); Vagrant (V) one record per several years (non-residents).

Excluding Vagrants, commoner species which were recorded in the area before 2002 but not seen between 2002 and 2013, including during this study, are regarded as “lost”, while those not recorded prior to 2002 are termed “gained”, even though some of them might have been present in the respective period but overlooked. Vagrants (as defined above) when lost or gained are represented by a V in either columns, and are not included in the totals of lost or gained. The bracket after the total lost or total gained gives the number of Vagrants excluded from the totals (Table 1).

A combination of wader nets (used for water birds) and mist nets (forest birds) with audio playback was used by TAA, TEA, TCO and GT (TAA was involved in all netting sessions) to trap birds at points shown on Fig. 2. Nets of varying lengths were used, ranging between 100 and 250 m and with height varying between 1.5 and 4 m. Netting sessions took place between 5h30 and 17h00, though sessions were shorter when the weather was not suitable for netting birds (rain or hot weather) or when the frequency of bird captures was low. Between five and 14 netting days were carried out each quarter, with rotation of nets to various habitats. Within each habitat type, nets were set up in different sites, and were moved after 2–7 days depending on trapping success. Net check interval was 20 min., to minimise heat stress and exhaustion of trapped birds. More netting time was spent in forest because of the greater probability of missing understorey birds during transect walks.

We collate and compare our 2009–13 transect and mist-netting data with earlier records, to reveal the trends in bird species composition over time. Taxonomic treatment follows Borrow & Demey (2001).

Results

The birds recorded at IITA in the present study, and in the Ibadan area by other recent and earlier studies, are summarized in Table 1.




Figure 2. Location of transects (thick black lines labelled Dam 1–3 and For 1–3) and mist-netting points (ovals) within the IITA campus IBA.

Table 1. Birds of the IITA campus IBA recorded in the present study, plus other recent and earlier records from IITA (sources underlined) and from the Ibadan area. Early records: Ban = Bannerman (1930–51); Wil = Willoughby (1949); S = Serle (1950); ES = Elgood & Sibley (1964); EFD = Elgood *et al.* (1973); EE/EE = Elgood *et al.* (1994); JB = Button (1965); DR = Robinson (1966); W = Wells (1966a, 1966b, 1967); B = Bass (1967); RP = Parker (1967, 1968, 1970); AP = Ashford & Parker (1968); P = Pettet (1968a, 1968b, 1975); A = Ashford (1968, 1969); NR = Robinson (1970); BB = Broadbent (1972); JA/JA = Ash (1990); AE = Ezealor (2001). Recent records: D = Demey *et al.* (2003); M = Manu *et al.* (2005); DB = D. Bengtsson (*in Demey* 2006); TEA = Adeyanju & Adeyanju (2012); JP = J. Peacock (pers. comm.); GH = G. Huk (pers. comm.); TH = Thibault *et al.* (2012); TAA = T.A. Adeyanju observations outside IITA. Records from P. Hall (PH) span a long period and each record is classed according to its precise date. “Gained” = species recorded in the Ibadan area since 2002 but not before; “lost” = species recorded before 2002 but not since. * = species restricted to Guinea–Congo Rain-forest biome; ** = species restricted to Sudan–Guinea Savanna biome (Ezealor 2001); † = species observed in this study within the forest reserve; ‡ = unconfirmed records.

| | Observed this study | Netted this study | Other recent records | Early records | Gained | Lost |
|--|------------------------|----------------------|---------------------------|----------------------|--------|------|
| Podicipitidae | | | | | | |
| <i>Tachybaptus ruficollis</i> Little Grebe | | | GH (2012) | ES (sporadic), B, BB | | |
| Phalacrocoracidae | | | | | | |
| <i>Phalacrocorax africanus</i> Long-tailed Cormorant | F ¹ | | | ES | | |
| Anhingidae | | | | | | |
| <i>Anhinga rufa</i> African Darter | | | | ES | | X |
| Ardeidae | | | | | | |
| <i>Botaurus stellaris</i> Great Bittern | | | | PH (2001), EE | | V |
| <i>Exobrychus minutus</i> Little Bittern | F | I | | ES, BB | | |
| <i>I. sturmi</i> Dwarf Bittern | | | | PH (1998), ES | | X |
| <i>Gorsachius leuconotus</i> White-backed Night Heron | | | | ES (once) | | V |
| <i>Nycticorax nycticorax</i> Black-crowned Night Heron | F ² | | | BB | | |
| <i>Ardeola ralloides</i> Squacco Heron | A | | | ES | | |
| <i>Bubulcus ibis</i> Cattle Egret | A | | TAA(2009–14) ³ | ES, B | | |

| | | | | |
|--|----------------|------------------------|---|---|
| <i>Butorides striata</i> Green-backed Heron | C | ES | | |
| <i>Egretta ardesiaca</i> Black Heron | | <u>PH</u> ⁴ | X | |
| <i>E. gularis</i> Western Reef Heron | | <u>EE</u> | V | |
| <i>E. garzetta</i> Little Egret | U | ES | | |
| <i>E. intermedia</i> Intermediate Egret | C | ES, EE | | |
| <i>E. alba</i> Great Egret | C | ES | | |
| <i>Ardea purpurea</i> Purple Heron† | C | ES | | |
| <i>A. cinerea</i> Grey Heron | C | ES | | |
| <i>A. melanocephala</i> Black-headed Heron | C | ES | | |
| Scopidae | | | | |
| <i>Scopus umbretta</i> Hamerkop | | ES, P | | |
| Ciconiidae | | | | |
| <i>Mycteria ibis</i> Yellow-billed Stork | | ES, DR | V | |
| <i>Anastomus lamelligerus</i> African Openbill Stork | | ES | V | |
| <i>Ciconia abdimii</i> Abdim's Stork | V | ES | | |
| <i>C. episcopus</i> Woolly-necked Stork | R | RP | | |
| <i>C. ciconia</i> White stork | | <u>PH (1998, 2000)</u> | V | |
| Threskiornithidae | | | | |
| <i>Plegadis falcinellus</i> Glossy Ibis | V ⁵ | | V | |
| <i>Bostrychia hagedash</i> Hadada Ibis | C ⁶ | | X | |
| <i>Threskiornis aethiopicus</i> Sacred Ibis | | ES | | V |

¹1–2 around the main lake, each year.²Included juveniles.³> 600 roost in UI Awba dam environs.⁴Three occasions 1993–5.⁵Once in 2010, a group of c. 6 birds.⁶Favour the small lake area beside the golf course.

| | Observed this study | Netted this study | Other recent records | Early records | Gained | Lost |
|---|------------------------|----------------------|-------------------------|------------------------------|--------|------|
| Anatidae | | | | | | |
| <i>Dendrocygna viduata</i> White-faced Whistling Duck | A ⁷ | | | Ban, EE | | |
| <i>Plectropterus gambensis</i> Spur-winged Goose | F ⁸ | | | PH, Wil | | |
| <i>Pteronetta hartlaubii</i> Hartlaub's Duck* | | | | PH (1995, a pair), AE | | V |
| <i>Sarkidiornis melanotos</i> Knob-billed Goose | | | | PH (1987, 1993, 1995), Wil | | V |
| <i>Nettapus auritus</i> African Pygmy Goose | | | | PH (until 1995), ES, ASH, EE | | X |
| <i>Anas penelope</i> Eurasian Wigeon | | | | Ban, EE | | V |
| <i>A. crecca</i> Common Teal | | | | PH (Jan 1994, a pair), EE | | V |
| <i>A. acuta</i> Northern Pintail | | | | PH (1989, 1998) | | V |
| <i>A. querquedula</i> Garganey | | | | PH (1993, 1994), Ban, EE | | V |
| <i>Aythya nyroca</i> Ferruginous Duck | | | | P | | V |
| Pandionidae | | | | | | |
| <i>Pandion haliaetus</i> Osprey† | U | | | ES, BB | | |
| Accipitridae | | | | | | |
| <i>Aviceda cuculoides</i> African Cuckoo Hawk† | U | | | ES, EE | | V |
| <i>Pernis ptilorivus</i> European Honey Buzzard† | U | | PH | | | |
| <i>Macheiramphus alcinus</i> Bat Hawk† | U ⁹ | | PH (May 2010) | | | |
| <i>Elanus caeruleus</i> Black-shouldered Kite | F | | | ES | | |
| <i>Milvus migrans</i> Yellow-billed Kite† | A | | | ES | | |
| <i>Gypohierax angolensis</i> Palm-nut Vulture† | F | | | ES, W | | |
| <i>Polyboroides typus</i> African Harrier Hawk† | F | | | ES | | |
| <i>Circus macrourus</i> Pallid Harrier | U | | | ES | | |
| <i>C. aeruginosus</i> Eurasian Marsh Harrier† | U | | | Ban | | V |
| <i>Micronisus gabar</i> Gabar Goshawk | F | | | BB | | |
| <i>Accipiter tachiro</i> African Goshawk† | C | 5 | | ES, NR | | X |
| <i>A. badius</i> Shikra | F | | | ES, B, NR | | |

| | | | | | | |
|---|------------------|---|--|-----------------|------------------------------------|---|
| <i>A. erythropus</i> Red-thighed Sparrowhawk**† | | | | <u>PH</u> | <u>AE</u> ES | X |
| <i>A. minullus</i> Little Sparrowhawk | | | | | <u>PH, ES</u> | X |
| <i>A. melanoleucus</i> Black Sparrowhawk | | | | | <u>AE</u> ES | V |
| <i>Urotriorchis macrourus</i> Long-tailed Hawk**† | | | | <u>PH, D</u> | ES | |
| <i>Kaupifalco monogrammicus</i> Lizard Buzzard† | C | I | | | EE | |
| <i>Buteo auguralis</i> Red-necked Buzzard† | F | | | | | |
| <i>Aquila rapax</i> Tawny Eagle | U | | | <u>PH (one)</u> | ES, BB | X |
| <i>Hieraaetus spilogaster</i> African Hawk-Eagle† | F | | | <u>PH</u> | ES, EE ES | |
| <i>Lophaetus occipitalis</i> Long-crested Eagle† | C | | | | ES, BB, NR ES | |
| Falconidae | | | | | | |
| <i>Falco tinnunculus</i> Common Kestrel | C | | | | ES, EE ES | |
| <i>F. ardosiaceus</i> Grey Kestrel† | F | | | | ES, BB, NR ES | |
| <i>F. cucullatus</i> African Hobby† | F | | | | | |
| <i>F. biarmicus</i> Lanner† | U ¹⁰ | | | | | |
| Phasianidae | | | | | | |
| <i>Coturnix delegorguei</i> Harlequin Quail | | | | | JB, EE | V |
| <i>Ptilopachus petrosus</i> Stone Partridge | | | | | ES (rare, Ojo Hillis) ES (once) | X |
| <i>Francolinus lathamii</i> Latham's Francolin* | | | | | <u>PH</u> ¹¹ | V |
| <i>F. squamatus</i> Scaly Francolin | C ¹² | | | | <u>AE, Ban</u> ES | V |
| <i>F. achantensis</i> Ahanta Francolin**† | VA ¹³ | | | | | |
| <i>F. bicalcaratus</i> Double-spurred Francolin† | | | | | | |

⁷Present all year; though numbers previously observed to increase in dry season, large flocks observed in rains of 2011 and 2012.

⁸Bred in 2009 but in 2010 and 2011 only one individual observed. PH observed them in 1995 but breeding not confirmed.

⁹A pair nested within the IBA, 2009–13.

¹⁰Pairs in the IBA on very tall trees (> 40 m).

¹¹Forest on the west bank of the main lake.

¹²Common at forest edges, only frequent in the interior.

¹³Mostly farmland but recently also present close to forest.

| | Observed this study | Netted this study | Other recent records | Early records | Gained | Lost |
|--|------------------------|----------------------|-------------------------|---------------------------------------|--------|------|
| Numididae | | | | | | |
| <i>Guttera pucherani</i> Crested Guineafowl | | | | EE | | V |
| <i>Numida meleagris</i> Helmeted Guineafowl | F ¹⁴ | | | | X | |
| Turnicidae | | | | | | |
| <i>Turnix sylvatica</i> Little Buttonquail | | | | ES (once) | | V |
| Rallidae | | | | | | |
| [<i>Sarothrura elegans</i> Buff-spotted Flufftail | | | DB (Oct 2005) | | V] | |
| <i>Crex egregia</i> African Crane | | U | | ES, W | | V |
| <i>Porzana porzana</i> Spotted Crane | | | | <u>PH (pair, Feb 1996)</u> | | X |
| <i>Aenigmatolimnas marginalis</i> Striped Crane | | C | | ES, W | | |
| <i>Amaurornis flavirostra</i> Black Crane | | C | | ES | | |
| <i>Porphyrio alleni</i> Allen's Gallinule | | C | | ES, EE | | |
| <i>P. porphyrio</i> Purple Swamphen | | | | <u>PH (1999, 2000)</u> | | V |
| <i>Gallinula chloropus</i> Common Moorhen | | C | | Ban | | |
| <i>G. angulata</i> Lesser Moorhen | | R | | ES | | |
| Helionithidae | | | | | | |
| <i>Podica senegalensis</i> African Finfoot | | | TEA | <u>PH (bred Aug 1999)</u> , ES, BB, W | | |
| Otididae | | | | | | |
| <i>Lissotis melanogaster</i> Black-bellied Bustard | | | | W, ES | | V |
| Jacaniidae | | | | | | |
| <i>Actophilornis africanus</i> African Jacana | VA ¹⁵ | 15 | | ES | | |
| Rostratulidae | | | | | | |
| <i>Rostratula benghalensis</i> Greater Painted-snipe | C ¹⁶ | 13 | | ES, <u>EE</u> | | |
| Recurvirostridae | | | | | | |
| <i>Himantopus himantopus</i> Black-winged Stilt | F | | | ES, BB | | |

| | | | | | | | | | |
|------------------------------|-------------------------|-------------------|---|----------------------------------|--|--|--|--|---|
| Burhinidae | | | | | | | | | |
| <i>Burhinus senegalensis</i> | Senegal Thick-knee | C ¹⁷ | | ES | | | | | |
| Glareolidae | | | | | | | | | |
| <i>Pluvianus aegyptius</i> | Egyptian Plover | V | | PH, ES, EE | | | | | |
| <i>Glareola pratincola</i> | Collared Pratincole | V (one, Jan 2011) | | PH (Sep 1994), ES, EE | | | | | V |
| <i>G. cinerea</i> | Grey Pratincole | | | | | | | | |
| Charadriidae | | | | | | | | | |
| <i>Charadrius dubius</i> | Little Ringed Plover | | | PH ¹⁸ , ES (frequent) | | | | | X |
| <i>C. hiaticula</i> | Common Ringed Plover | | | PH (one, Sep 1994), ES (rare) | | | | | V |
| <i>C. forbesi</i> | Forbes's Plover | F | | ES, EE | | | | | |
| <i>Vanellus senegallus</i> | African Wattled Lapwing | V | | | | | | | V |
| <i>V. albiceps</i> | White-headed Lapwing | C | | ES | | | | | |
| <i>V. spinosus</i> | Spur-winged Lapwing | A | 5 | EE | | | | | |
| Scolopacidae | | | | | | | | | |
| <i>Calidris temminckii</i> | Temminck's Stint | | | | | | | | V |
| <i>C. ferruginea</i> | Curlew Sandpiper | | | | | | | | V |
| <i>Philomachus pugnax</i> | Ruff | | | | | | | | V |
| <i>Lymnocyptes minimus</i> | Jack Snipe | U | | | | | | | X |
| <i>Gallinago gallinago</i> | Common Snipe | U | 2 | ES, P, BB, EE | | | | | X |
| <i>G. media</i> | Great Snipe | U | | | | | | | |
| <i>Numenius arquata</i> | Eurasian Curlew | | | | | | | | V |
| <i>Tringa erythropus</i> | Spotted Redshank | | | | | | | | |
| <i>T. totanus</i> | Common Redshank | | | | | | | | V |

¹⁴Juveniles observed. Not recorded within the forest reserve, but feathers often found around cassava plots in IITA fields.

¹⁵Juveniles in early rains.

¹⁶A few resident all year and breeding. ES considered it a migrant, present Mar–Jun.

¹⁷In mixed flocks of plovers at the IITA main lake.

¹⁸A few present during the dry season every year 1987–95.

| | Observed this study | Netted this study | Other recent records | Early records | Gained Lost |
|--|------------------------|----------------------|-------------------------|---|-------------|
| <i>T. stagnatilis</i> Marsh Sandpiper | V | | | <u>PH</u> (two, Oct 1994) | V |
| <i>T. nebularia</i> Common Greenshank | | | | W | |
| <i>T. ochropus</i> Green Sandpiper | | | <u>PH</u> | ES | |
| <i>T. glareola</i> Wood Sandpiper | F | | | ES | |
| <i>T. hypoleucos</i> Common Sandpiper | C ¹⁹ | | | ES | |
| <i>Arenaria interpres</i> Ruddy Turnstone | | | | <u>PH</u> (one, Sep 1991) | V |
| Laridae | | | | | |
| <i>Larus ridibundus</i> Black-headed Gull | | | D | | V |
| Sternidae | | | | | |
| <i>Gelochelidon nilotica</i> Gull-billed Tern | | | | <u>PH</u> (one, Oct 1994) | V |
| <i>Chlidonias leucopterus</i> White-winged Tern | | | | <u>PH</u> (1988, 1989, 2000), <u>EE</u> | V |
| Rhynchopidae | | | | | |
| <i>Rynchops flavisrostris</i> African Skimmer | | | | <u>EE</u> (one) | V |
| Columbidae | | | | | |
| <i>Treeron calvus</i> African Green Pigeon† | C | | | ES | |
| <i>Turtur brehmeri</i> Blue-headed Wood Dove**† | | | <u>PH</u> ²⁰ | <u>AE</u> | |
| <i>T. tympanistris</i> Tambourine Dove† | C | 2 | | ES | |
| <i>T. afer</i> Blue-spotted Wood Dove† | A | | | ES, NR | X |
| <i>Columba iriditorques</i> Western Bronze-naped Pigeon* | | | | <u>PH</u> ²¹ | |
| <i>C. guinea</i> Speckled Pigeon | C | | | <u>EE</u> ²² | X |
| <i>Streptopelia semitorquata</i> Red-eyed Dove† | A ²³ | 1 | | ES | |
| <i>S. vinacea</i> Vinaceous Dove | R ²⁴ | | <u>PH</u> | | X |
| <i>S. senegalensis</i> Laughing Dove | C ²⁵ | | | ES, NR | |
| Pittidae | | | | | |
| <i>Poicephalus senegalus</i> Senegal Parrot**† | R | | TAA | <u>AE</u> , ES, <u>EE</u> | X |
| <i>Agapornis pullarius</i> Red-headed Lovebird | | | | BB | |

| | | | | | | | | |
|------------------------------|------------------------------|-----------------|-------------------------|------------------------------------|--|--|--|---|
| Musophagidae | | | | | | | | |
| <i>Tauraco persa</i> | Green Tauraco*† | C | <u>M</u> | <u>AE, ES</u> | | | | V |
| <i>Musophaea violacea</i> | Violet Tauraco** | C | | <u>AE, P</u> ES, JA | | | | |
| <i>Crinifer piscator</i> | Western Grey Plantain-eater† | | | | | | | |
| Cuculidae | | | | | | | | |
| <i>Oxyophus jacobinus</i> | Jacobin Cuckoo† | U | | | | | | X |
| <i>O. levaillantii</i> | Levaillant's Cuckoo† | U | <u>M</u> | ES, B | | | | |
| <i>Clamator glandarius</i> | Great Spotted Cuckoo | U | | ES, B, P | | | | X |
| <i>Cuculus solitarius</i> | Red-chested Cuckoo | | <u>PH (few. rains)</u> | | | | | X |
| <i>C. clamosus</i> | Black Cuckoo† | U | | | | | | |
| <i>C. canorus</i> | Common Cuckoo | | | | | | | |
| <i>C. gularis</i> | African Cuckoo† | | <u>PH</u> ²⁶ | <u>PH (one. May 1998)</u> W, BB | | | | V |
| <i>Chrysococcyx cupreus</i> | African Emerald Cuckoo† | U | <u>M</u> | ES | | | | |
| <i>C. klaas</i> | Klaas's Cuckoo† | F | | ES, NR | | | | |
| <i>C. caprius</i> | Didric Cuckoo† | F | | ES, NR | | | | |
| <i>Ceuthochares aereus</i> | Yellowbill† | C | <u>M</u> | ES | | | | |
| <i>Centropus leucogaster</i> | Black-throated Coucal* | R | | <u>AE, ES</u> | | | | |
| <i>C. grillii</i> | Black Coucal† | U | | | | | | X |
| <i>C. senegalensis</i> | Senegal Coucal† | A ²⁷ | | ES, NR | | | | |

¹⁹Present throughout the year; numbers increase during dry season.

²⁰Infrequent on west bank of main lake.

²¹Uncommonly heard during rains in 2000.

²²First record Ibadan area 1960.

²³Observed in all habitat types present.

²⁴A savanna species, now occasionally sighted in IITA and environs.

²⁵Only near residences.

²⁶Occasional throughout dry season.

²⁷Black variant "epomidis". Frequent.

| | Observed this study | Netted this study | Other recent records | Early records | Gained Lost |
|---|------------------------|----------------------|-------------------------|---------------------------------|----------------|
| <i>C. monachus</i> Blue-headed Coucal† | F | | | ES | |
| Tytonidae | | | | | |
| <i>Tyto alba</i> Barn Owl | U | | PH | ES | |
| Strigidae | | | | | |
| <i>Otus scops</i> European Scops Owl | | | | ES, W, BB, EE | X |
| <i>Ptilopsis leucotis</i> Northern White-faced Owl† | C ²⁸ | 1 | | ES | |
| <i>Bubo africanus</i> Spotted Eagle Owl | | | | ES | X |
| <i>B. poensis</i> Fraser's Eagle Owl*† | C (2011) | | TAA | ES, EE PH ²⁹ , ES | |
| <i>Glaucidium perlatum</i> Pearl-spotted Owlet | | | | ES | |
| <i>Strix woodfordii</i> African Wood Owl | C | | | | |
| Caprimulgidae | | | | | |
| <i>Caprimulgus climacurus</i> Long-tailed Nightjar | C | | PH ³⁰ | ES, BB | |
| <i>C. nigriscapularis</i> Black-shouldered Nightjar* | | | PH (dry season) | AE, Ban, ES | |
| <i>C. inornatus</i> Plain Nightjar | | | | ES | |
| <i>C. europaeus</i> European Nightjar | (once, 2010) | | | EE | |
| <i>Macrodipteryx longipennis</i> Standard-winged Nightjar | U | | | ES, BB | X |
| <i>M. vexillarius</i> Pennant-winged Nightjar | | | | RP | |
| Apodidae | | | | | |
| <i>Talacanthura ussheri</i> Mottled Spinetail† | C | | | P | X |
| <i>Cypsiurus parvus</i> African Palm Swift | C | | | ES | X |
| <i>Apus pallidus</i> Pallid Swift | | | | P | X |
| <i>A. apus</i> Common Swift | | | | PH (flock of 10), ES | X |
| <i>A. caffer</i> White-rumped Swift | | | | PH ³¹ | |
| <i>A. affinis</i> Little Swift | U | | | ES | |
| Alcedinidae | | | | | |
| <i>Halcyon leucocephala</i> Grey-headed Kingfisher | R | | | ES | |

| | | | | | | | | |
|--|--------|---|--|-----------------------------|--|-----------|--|----|
| <i>H. malimbica</i> Blue-breasted Kingfisher† | C | 2 | | ES, NR | | | | |
| <i>H. senegalensis</i> Woodland Kingfisher† | A | 5 | | ES, NR | | | | X |
| <i>H. chelicuti</i> Striped Kingfisher | | | | ES | | | | |
| <i>Ceyx tecontei</i> African Dwarf Kingfisher | (once) | | | JA, EE | | | | |
| <i>C. pictus</i> African Pygmy Kingfisher | R | 1 | | ES, NR | | | | |
| <i>Alcedo cristata</i> Malachite kingfisher | C | 3 | | ES | | | | |
| <i>Megaceryle maxima</i> Giant Kingfisher | R | | | ES | | | | |
| <i>Ceryle rudis</i> Pied Kingfisher | F | | | ES | | | | |
| Meropidae | | | | | | | | |
| <i>Merops pusillus</i> Little Bee-eater | | | | ES | | | | V |
| <i>M. albicollis</i> White-throated Bee-eater† | C | | | ES, NR | | | | V |
| <i>M. apiaster</i> European Bee-eater | | | | P, EE | | | | |
| <i>M. malimbicus</i> Rosy Bee-eater*† | F | | | <u>AE, ES, P</u> | | <u>PH</u> | | V |
| <i>M. nubicus</i> Northern Carmine Bee-eater | | | | ES | | | | |
| Coraciidae | | | | | | | | |
| <i>Coracias abyssinicus</i> Abyssinian Roller | | | | ES (once), BB | | | | V |
| [<i>C. garrulus</i> European Roller | | | | ES (rare), P, BB (possible) | | | | V] |
| <i>Eurystomus gularis</i> Blue-throated Roller*† | R | | | <u>AE, ES</u> | | | | |
| <i>E. glaucurus</i> Broad-billed Roller† | F | | | <u>EE</u> | | <u>M</u> | | |
| Phoeniculidae | | | | | | | | |
| <i>Phoeniculus boilei</i> White-headed Wood-hoopoe | | | | ES, EE | | | | X |
| <i>P. purpureus</i> Green Wood-hoopoe† | F | | | ES, NR, EE | | | | X |
| <i>Rhinopomastus aterrimus</i> Black Wood-hoopoe | | | | W, EE | | | | |

²⁸Three young collected in Jan 2014.

²⁹Last observed in IITA in 2000, a pair calling throughout the rains.

³⁰Throughout dry season, restricted to area on west bank of main lake.

³¹A pair breeding, Oct 1990.

| | Observed this study | Netted this study | Other recent records | Early records | Gained Lost |
|--|------------------------|----------------------|-------------------------|------------------------------|-----------------|
| Bucerotidae | | | | | |
| <i>Tropicranus albo cristatus</i> White-crested Hornbill*† | U | | <u>M</u> | <u>AE</u> , ES, S | X |
| <i>Tockus camurus</i> Red-billed Dwarf Hornbill | | | | ES | |
| <i>T. fasciatus</i> African Pied Hornbill*† | A | | | <u>AE</u> , ES | |
| <i>T. nasutus</i> African Grey Hornbill† | C | | | ES | |
| <i>Bycanistes fistulator</i> Piping Hornbill | | | | <u>PH</u> ³² , ES | X |
| Capitonidae | | | | | |
| <i>Gymnobucco peii</i> Bristle-nosed Barbet* | | | | <u>AE</u> | X |
| <i>G. calvus</i> Naked-faced Barbet*† | | | <u>M</u> | <u>PH</u> , <u>AE</u> , ES | X ³³ |
| <i>Pogonius scolopaceus</i> Speckled Tinkerbird*† | C | 3 | | <u>AE</u> , <u>EE</u> | |
| <i>P. atroflavus</i> Red-rumped Tinkerbird*† | C | | | EE | |
| <i>P. subsulphureus</i> Yellow-throated Tinkerbird*† | C | | | <u>AE</u> , ES | |
| <i>P. bilineatus</i> Yellow-rumped Tinkerbird† | C | | | | X |
| <i>P. chrysoconus</i> Yellow-fronted Tinkerbird | | | | EE | |
| <i>Tricholaema hirsuta</i> Hairy-breasted Barbet*† | F | | <u>M</u> | <u>AE</u> , ES, EE | X |
| <i>Lybius vieilloti</i> Vieillot's Barbet† | U | | | ES, EE | |
| <i>L. bidentatus</i> Double-toothed Barbet | | | TAA, TEA | ES, EE | |
| <i>Trachyphonus purpuratus</i> Yellow-billed Barbet*† | U | | | <u>AE</u> , ES | |
| Indicatoridae | | | | | |
| <i>Prodotiscus insignis</i> Cassin's Honeyguide* | | | | ES | X |
| <i>Melichneutes robustus</i> Lyre-tailed Honeyguide* | | | | <u>AE</u> | X |
| <i>Indicator maculatus</i> Spotted Honeyguide*† | C | 1 | | | X |
| <i>I. indicator</i> Greater Honeyguide | | | | ES, EE | |
| <i>I. minor</i> Lesser Honeyguide | | | TAA (2013) | <u>PH</u> ³⁴ , ES | |
| <i>I. exilis</i> Least Honeyguide | | | | ES | X |

| | | | | | | | | | |
|---------------------------------|---------------------------|---|---|--|--|--|--|---|----|
| Picidae | | | | | | | | | |
| <i>Campethera punctuligera</i> | Fine-spotted Woodpecker | | | | | | | | X |
| <i>C. cailliautii</i> | Green-backed Woodpecker | | | | | | | PH (one, May 1998) | V |
| <i>C. nivosa</i> | Buff-spotted Woodpecker*† | F | 6 | | | | | ES (rare), EE PH ³⁵ , AE | V |
| <i>Dendropicus gabonensis</i> | Gabon Woodpecker* | | | | | | | EE | X |
| <i>D. fuscescens</i> | Cardinal woodpecker | | | | | | | AE, ES, EE | |
| <i>D. pyrrhogaster</i> | Fire-bellied Woodpecker*† | F | 2 | | | | | PH TAA | |
| <i>D. goertae</i> | Grey Woodpecker | | | | | | | PH (1987, 1988), ES | |
| Eurylaemidae | | | | | | | | | |
| <i>Smithornis rufolateralis</i> | Rufous-sided Broadbill*† | F | 1 | | | | | | X |
| Pittidae | | | | | | | | | |
| <i>Pitta angolensis</i> | African Pitta | | | | | | | W | X |
| Hirundinidae | | | | | | | | | |
| [<i>Riparia paludicola</i>] | Plain Martin | | | | | | | A, EE | VJ |
| <i>R. riparia</i> | Common Sand Martin | F | | | | | | TAA(2013) PH(Mar 1992, Oct 1994), BB, A, EE | VJ |
| [<i>R. cincta</i>] | Banded Martin | | | | | | | A (one) | |
| <i>Hirundo semirufa</i> | Rufous-chested Swallow | F | | | | | | ES, NR, A, EFD | X |
| <i>H. senegalensis</i> | Mosque Swallow | | | | | | | PH(frequent all year), ES, A, EFD, EE | |
| <i>H. abyssinica</i> | Lesser Striped Swallow | C | | | | | | ES, A, EFD | |
| <i>H. fuligula</i> | Rock Martin | F | | | | | | | X |
| <i>H. smithii</i> | Wire-tailed Swallow | | | | | | | ES | X |
| <i>H. leucosoma</i> | Pied-winged Swallow* | | | | | | | AE, ES, EE | X |
| <i>H. aethiopica</i> | Ethiopian Swallow | C | | | | | | ES, A | |

³²A few, Jan–Mar 1999 and Oct 2000.

³³Recorded by ES as frequent and by PH as a common breeder in the 1990s. Although M found it, seems now to have disappeared from the campus.

³⁴A pair parasitizing Naked-faced Barbet nests, Feb 1995.

³⁵A pair feeding chicks, Aug 1999.

| | Observed this study | Netted this study | Other recent records | Early records | Gained Lost |
|---|------------------------|----------------------|-------------------------|-------------------------------------|----------------|
| <i>H. rustica</i> Barn Swallow | F | | | ES, A ES | X |
| <i>Delichon urbicum</i> House Martin | | | | | |
| Motacillidae | | | | | |
| <i>Motacilla flava</i> Yellow Wagtail | F | | | ES, B, A, NR ES, NR | |
| <i>M. aguimp</i> African Pied Wagtail | C | | | | |
| <i>Anthus leucophrys</i> Plain-backed Pipit | C | | TH (2012) | ES, EFD | V |
| <i>A. pallidiventris</i> Long-legged Pipit | | | | | |
| <i>A. irivialis</i> Tree Pipit | | | | PH, P, BB PH, P, BB, A ES, EE | X X |
| <i>A. cervinus</i> Red-throated Pipit | F | | | | |
| <i>Macronyx croceus</i> Yellow-throated Longclaw | | | | | |
| Campephagidae | | | | | |
| <i>Campephaga phoenicea</i> Red-shouldered Cuckoo-shrike† | U | | M | ES, B, BB, EFD AE | X |
| <i>Coracina azurea</i> Blue Cuckoo-shrike* | | | | | |
| Pycnonotidae | | | | | |
| <i>Andropadus virens</i> Little Greenbul*† | A | 57 | | ES | |
| <i>A. gracilis</i> Little Grey Greenbul† | U | 2 | M | ES | |
| <i>A. curvirostris</i> Cameroon Sombre Greenbul*† | F | 15 | | AE | |
| <i>A. gracilirostris</i> Slender-billed Greenbul† | C | | | ES (rare), EE | |
| <i>A. latirostris</i> Yellow-whiskered Greenbul† | A | 109 | | ES | |
| <i>Baeopogon indicator</i> Honeyguide Greenbul*† | F | 4 | | AE, ES AE, EE | X |
| <i>Ixonotus guttatus</i> Spotted Greenbul* | C | 1 | | AE, ES AE, ES | |
| <i>Chlorocichla simplex</i> Simple Leaflove*† | U ³⁶ | | | | |
| <i>C. flavicollis</i> Yellow-throated Leaflove† | C | | | AE, ES | |
| <i>Thestelocichla leucopleura</i> Swamp-palm Bulbul*† | C | 1 | | AE, ES AE, ES | |
| <i>Pyrrhurus scandens</i> Leaflove*† | C | | | AE, ES | |
| <i>Phyllastrephus baumanni</i> Baumann's Greenbul*† | U | 10 | | AE, ES | |

| | | | | | | | |
|-----------------------------------|-------------------------------|---|----|----------|--|---------------------------|---|
| <i>P. icterinus</i> | Icterine Greenbul* | | | | | <u>AE</u> | X |
| <i>P. albicularis</i> | White-throated Greenbul**† | F | 26 | | | <u>AE</u> , ES | X |
| <i>Bleda syndactylus</i> | Red-tailed Bristlebill† | | | | | P, EE | |
| <i>B. canicapillus</i> | Grey-headed Bristlebill**† | C | 87 | | | <u>AE</u> , ES | |
| <i>Crinifer barbatus</i> | Western Bearded Greenbul**† | | | <u>M</u> | | Ban, <u>AE</u> , EE | |
| <i>C. calurus</i> | Red-tailed Greenbul* | U | | <u>M</u> | | <u>AE</u> , ES | X |
| <i>C. ndussumensis</i> | White-bearded Greenbul* | | | | | <u>AE</u> | |
| <i>Pycnonotus barbatus</i> | Common Bulbul† | A | 3 | <u>M</u> | | ES, NR | |
| <i>Nicator chloris</i> | Western Nicator**† | F | 7 | | | ES | |
| Turdidae | | | | | | | |
| <i>Stiphrornis erythrothorax</i> | Forest Robin**† | C | 15 | | | <u>AE</u> , ES (rare), BB | |
| <i>Luscinia megarhynchos</i> | Common Nightingale | | | | | ES, P, A, EE | |
| <i>L. svecica</i> | Bluethroat | | | | | A | V |
| <i>Cossypha cyanocamptor</i> | Blue-shouldered Robin Chat**† | F | 5 | | | Ban, EE | |
| <i>C. niveicapilla</i> | Snowy-crowned Robin Chat† | C | 2 | | | ES, NR, EFD | |
| <i>Neocossyphus poensis</i> | White-tailed Ant Thrush† | F | 4 | | | | X |
| <i>Stizorhina finschi</i> | Finsch's Flycatcher Thrush | | | | | <u>PH</u> , ES | X |
| <i>Phoenicurus phoenicurus</i> | Common Redstart | | | | | ES, A | X |
| <i>Saxicola rubetra</i> | Whinchat | C | | | | ES, A | V |
| <i>Oenanthe oenanthe</i> | Northern Wheatear | | | | | ES | V |
| <i>Monticola solitarius</i> | Blue Rock Thrush | | | | | W | V |
| <i>Turdus pelios</i> | African Thrush† | A | 5 | | | ES, NR, EFD | |
| Sylviidae | | | | | | | |
| <i>Melocichla mentalis</i> | African Moustached Warbler | F | | | | ES | X |
| <i>Acrocephalus schoenobaenus</i> | Sedge Warbler | | | | | A, BB | V |
| <i>A. scirpaceus</i> | European Reed Warbler | | | | | EE | V |
| <i>A. baeticatus</i> | African Reed Warbler | | | | | AP, A, BB, EE | X |

³⁶Savanna species termed common in Ibadan area by ES, now uncommon there; one pair observed in 2011, in farm bush by main lake.

| | Observed this study | Netted this study | Other recent records | Early records | Gained Lost |
|--|------------------------|----------------------|-------------------------|------------------|-------------|
| <i>A. arundinaceus</i> Great Reed Warbler | F | 2 | | ES, AP, A, BB | |
| <i>Hippolais polyglotta</i> Melodious Warbler | | | PH | ES, B, A, BB | |
| <i>H. icterina</i> Icterine Warbler | | | | RP, BB | X |
| <i>Cisticola erythropus</i> Red-faced Cisticola | A | 2 | | ES, A | X |
| <i>C. lateralis</i> Whistling Cisticola | | | | PH, ES | X |
| <i>C. anonymus</i> Chattering Cisticola* | | | | AE | X |
| <i>C. galactotes</i> Winding Cisticola | | | | PH | X |
| <i>C. brachypterus</i> Short-winged Cisticola | F | | | ES, A | |
| <i>C. juncidis</i> Zitting Cisticola | | | | PH | X |
| <i>Prinia subflava</i> Tawny-flanked Prinia | F | | | PH, ES | |
| <i>Heliotais erythropterus</i> Red-winged Warbler | | | | ES | X |
| <i>Camaroptera brachyura</i> Grey-backed Camaroptera | A | 7 | D | ES, S | |
| <i>C. supercilialis</i> Yellow-browed Camaroptera*† | C | 2 | | AE, ES | |
| <i>C. chloronota</i> Olive-green Camaroptera*† | A | 49 | D | ES | |
| <i>Macrosphenus kempii</i> Kemp's Longbill*† | F | 5 | | | X |
| <i>M. concolor</i> Grey Longbill*† | U | 2 | | ES | |
| <i>Eremomela pusilla</i> Senegal Eremomela** | U | | | AE, ES | |
| <i>Sylvietta brachyura</i> Northern Crombce | | | | EE | |
| <i>S. virens</i> Green Crombec*† | C | 3 | | AE, ES | V |
| <i>S. denti</i> Lemon-bellied Crombec*† | F | | | | |
| <i>Phylloscopus trochilus</i> Willow Warbler | F | | PH | ES, A, NR | X |
| <i>P. sibilatrix</i> Wood Warbler† | F | | | ES | |
| <i>Hypergerus atriceps</i> Oriole Warbler**† | F | | | AE, ES | |
| <i>Sylvia borin</i> Garden Warbler | U | | | ES, A, B, RP | |
| <i>S. communis</i> Common Whitethroat | | | | BB | V |
| <i>Hylia prasina</i> Green Hylia*† | C | 9 | | AE, ES | |

| | | | | | | | | | |
|------------------------------|------------------------------------|---------------|----|--|--|--|--|--|---|
| Muscicapidae | | | | | | | | | |
| <i>Fraseria ocreata</i> | Fraser's Forest Flycatcher**† | | | | | | | | X |
| <i>Melaneris edoloides</i> | Northern Black Flycatcher† | U (F in 2009) | | | | | | | |
| <i>Muscicapa striata</i> | Spotted Flycatcher | (once) | | | | | | | X |
| <i>M. olivaceus</i> | Olivaceous Flycatcher**† | U | 1 | | | | | | |
| <i>M. comitata</i> | Dusky-blue Flycatcher**† | | | | | | | | |
| <i>M. tessmanni</i> | Tessmann's Flycatcher** | | | | | | | | |
| <i>Ficedula hypoleuca</i> | Pied Flycatcher | | | | | | | | |
| Monarchidae | | | | | | | | | |
| <i>Erythrocerus mecallii</i> | Chestnut-capped Flycatcher** | | | | | | | | |
| <i>Trochocercus nitens</i> | Blue-headed Crested Flycatcher**† | F | 1 | | | | | | |
| <i>Terpsiphone viridis</i> | African Paradise Flycatcher† | | | | | | | | |
| <i>T. rufiventer</i> | Red-bellied Paradise Flycatcher**† | A | 41 | | | | | | |
| Platyteiridae | | | | | | | | | |
| <i>Megabyas flammulatus</i> | Shrike Flycatcher* | | | | | | | | |
| <i>Bias musicus</i> | Black-and-White Flycatcher | | | | | | | | |
| <i>Dyaphorophya castanea</i> | Chestnut Wattle-eye**† | F | 1 | | | | | | |
| <i>D. blissetti</i> | Red-cheeked Wattle-eye**† | C | 21 | | | | | | |
| <i>Platysteira cyanea</i> | Common Wattle-eye† | F | 1 | | | | | | |
| <i>Batis senegalensis</i> | Senegal Batis | | | | | | | | |
| Timaliidae | | | | | | | | | |
| <i>Illadopsis rufipennis</i> | Pale-breasted Illadopsis† | U | | | | | | | |
| <i>I. fulvescens</i> | Brown Illadopsis**† | C | 14 | | | | | | |
| <i>I. puveli</i> | Puvel's Illadopsis | | | | | | | | |
| <i>Turdoides plebejus</i> | Brown Babbler | | | | | | | | |

³⁷Observed on several occasions in mixed parties.

³⁸PH regarded it as common in mixed parties, more often heard than seen, and its call may initially have been confused with that of *I. fulvescens* which is common on the site.

| | Observed this study | Netted this study | Other recent records | Early records | Gained Lost |
|--|------------------------|----------------------|------------------------------------|--|----------------|
| <i>T. reinwardtii</i> Blackcap Babbler | F | 4 | | ES <u>AE</u> , ES | X |
| <i>Phyllanthus atripennis</i> Capuchin Babbler**† | | | | | |
| Remizidae | | | | | |
| <i>Pholidornis rufiae</i> Tit-hyllia | | | <u>PH</u> (several, <u>W</u> bank) | | X |
| Nectariniidae | | | | | |
| <i>Anthreptes rectirostris</i> Green Sunbird* | | | | Ban ES, NR | X |
| <i>Cyanomitra verticalis</i> Green-headed Sunbird | U | | | <u>PH</u> , <u>AE</u> , ES, B | |
| <i>C. cyanolaema</i> Blue-throated Brown Sunbird* | U ³⁹ | 1 | | ES | |
| <i>C. obscura</i> Western Olive Sunbird† | C | 75 | | <u>AE</u> , ES | |
| <i>Chalcomitra adelberti</i> Buff-throated Sunbird**† | U | | | | X |
| <i>C. senegalensis</i> Scarlet-chested Sunbird | R | | | | |
| <i>Hedydipna collaris</i> Collared Sunbird† | A | 32 | <u>M</u> | ES | |
| <i>Cinnyris chloropygius</i> Olive-bellied Sunbird† | F | 1 | | ES, NR | X |
| <i>C. minullus</i> Tiny Sunbird**† | U | 5 | | | |
| <i>C. venustus</i> Variable Sunbird | F | | | ES, NR, EFD <u>AE</u> , ES | X |
| <i>C. superbus</i> Superb sunbird* | | | | <u>AE</u> , ES, EFD | |
| <i>C. coccinigerastus</i> Splendid Sunbird*† | C | 5 | | ES, NR | |
| <i>C. cupreus</i> Copper Sunbird | C | 2 | | | |
| Zosteropidae | | | | | |
| <i>Zosterops senegalensis</i> Yellow White-eye | | | | <u>PH</u> & ES (common), NR | X |
| Laniidae | | | | | |
| <i>Lanius senator</i> Woodchat Shrike | | | | <u>PH</u> , ES | X |
| <i>Corvinella corvina</i> Yellow-billed Shrike* | F | | | <u>AE</u> , ES, A, <u>EE</u> | |
| Malconotidae | | | | | |
| <i>Malconotus cruentus</i> Fiery-breasted Bush-Shrike**† (one group) | U | | <u>PH</u> (common), <u>D</u> | <u>AE</u> , ES, B, NR ES, NR, <u>EE</u> | |
| <i>M. multicolor</i> Many-coloured Bush-Shrike† | | | | | |

| | | | | | |
|---|---|------------|--------------------|--------------------|---|
| <i>M. sulfurepectus</i> Sulphur-breasted Bush-shrike† | | | | ES, EFD | |
| <i>Antichromis minutus</i> Marsh Tchagra | | | | ES, EE | X |
| <i>Tchagra senegalus</i> Black-crowned Tchagra | | | | ES, B | X |
| <i>Dryocopus sabini</i> Sabine's Puffback*† | U | <u>M</u> | | <u>AE</u> , ES | |
| <i>D. gambensis</i> Northern Puffback | | TAA (2013) | | ES, NR | |
| <i>Laniarus aethiopicus</i> Tropical Boubout† | F | | | <u>EE</u> | |
| Prionopidae | | | | | |
| <i>Prionops plumatus</i> White Helmet-shrike | | | | ES | X |
| <i>P. caniceps</i> Red-billed Helmet-shrike*† | U | <u>M</u> | | <u>AE</u> , EE | |
| Oriolidae | | | | | |
| <i>Oriolus nigripennis</i> Black-winged Oriole*† | F | <u>M</u> | | <u>AE</u> , ES | |
| <i>O. brachyrhynchus</i> Western Black-headed Oriole*† | F | <u>M</u> | | | |
| <i>O. auratus</i> African Golden Oriole | | | PH (fairly common) | ES, EFD | |
| <i>O. oriolus</i> Eurasian Golden Oriole | | | | JA | V |
| Dicruridae | | | | | |
| <i>Dicrurus ludwigii</i> Square-tailed Drongo† | F | 2 | | ES | X |
| <i>D. atripennis</i> Shining Drongo*† | F | | | JA | X |
| <i>D. adsimilis</i> Fork-tailed Drongo | | | | ES, NR | |
| <i>D. modestus</i> Velvet-mantled Drongo† ⁴⁰ | C | 1 | <u>M</u> | | |
| Corvidae | | | | | |
| <i>Corvus albus</i> Pied Crow | A | | TAA | ES, EFD | V |
| <i>Psittosomus afer</i> Piapiac** | | | | | |
| Sturnidae | | | | | |
| <i>Poeoptera lugubris</i> Narrow-tailed Starling† | F | | | | X |
| <i>Orychognathus fulgidus</i> Forest Chestnut-winged Starling*† | F | | | <u>AE</u> , ES, NR | |

³⁹PH recorded it as fairly common on west bank of main lake, though seems much less common now; one juvenile trapped.

⁴⁰Many records were originally wrongly referred to the savanna species *D. adsimilis*, whereas the forest records should refer to *D. modestus*.

| | Observed this study | Netted this study | Other recent records | Early records | Gained Lost |
|---|------------------------|----------------------|------------------------------|---------------------------------|----------------|
| <i>Lamprolornis purpureus</i> Purple Glossy Starling | | | | ES | X |
| <i>L. splendidus</i> Splendid Glossy Starling | C | | | ES | |
| <i>L. caudatus</i> Long-tailed Glossy Starling | V (2009) | | | | V |
| <i>Cinnyricinclus leucogaster</i> Violet-backed Starling | | | | PH (dry season 1999), ES (once) | V |
| Passeridae | | | | | |
| <i>Passer griseus</i> Northern Grey-headed Sparrow | C | | | ES, NR | |
| Ploceidae | | | | | |
| <i>Plocepasser superciliosus</i> Chestnut-crowned Sparrow-Weaver* | | | | Ban | X |
| <i>Ploceus nigricollis</i> Black-necked Weaver | C | 38 | | NR | |
| <i>P. nigerrimus</i> Vieillot's Black Weaver* | U | | | AE | |
| <i>P. cucullatus</i> Village Weaver† | VA | 50 | | ES | |
| <i>P. tricolor</i> Yellow-mantled Weaver*† | C | | D, M | AE, ES | |
| <i>Malimbus nitens</i> Blue-billed Malimbe*† | A | 17 | M | AE, ES | |
| <i>M. malimbicus</i> Crested Malimbe*† | | | PH (frequent), M, TAA (2013) | AE, ES | |
| <i>M. ibadanensis</i> Ibadan Malimbe*† | R | | M | PH, AE, ES, JA | |
| <i>M. scutatus</i> Red-vented Malimbe*† | C | | M | AE, ES, NR | |
| <i>M. rubricollis</i> Red-headed Malimbe*† | C | | M | AE, ES, B | |
| <i>Quelea erythrops</i> Red-headed Quelea | F | 10 | | ES, BB | |
| <i>Euplectes hordeaceus</i> Black-winged Bishop | | | | Ban | X |
| <i>E. franciscanus</i> Northern Red Bishop | R | | | EE | |
| <i>E. macroura</i> Yellow-mantled Widowbird | U | | | ES, EFD, EE | |
| <i>Amblyospiza albifrons</i> Grosbeak Weaver | R | | | ES, B, BB | |
| Estrildidae | | | | | |
| <i>Nigrita canicapillus</i> Grey-crowned Negrofinch† | F | | M | ES | |
| <i>N. luteifrons</i> Pale-fronted Negrofinch* | | | | ES | |
| <i>N. bicolor</i> Chestnut-breasted Negrofinch*† | F | 1 | | AE, ES | X |

| | | | | | | |
|--|-----|----|----|--|--------------------|---------------|
| <i>N. fusconotus</i> White-breasted Nigrofinch**† | F | | | | | X |
| <i>Pyrenestes ostrinus</i> Black-bellied Seedcracker | U | I | | | ES, EE | |
| <i>Spermophaga haematina</i> Western Bluebill**† | C | 38 | | | <u>AE</u> , ES, BB | |
| <i>Mandingoa nitidula</i> Green Twinspot | | | | | Ban | X |
| <i>Lagonosticta rufopicta</i> Bar-breasted Firefinch**† | U | | | | AE, ES | |
| <i>L. rubricata</i> Blue-billed Firefinch | | | | | PH (frequent) | X |
| <i>Estrilda melpoda</i> Orange-cheeked Waxbill† | A | | | | ES, B, BB, EFD | |
| <i>Spermestes cucullatus</i> Bronze Mannikin† | C | 10 | | | ES | |
| <i>S. bicolor</i> Black-and-White Mannikin† | F | | | | ES | |
| Viduidae | | | | | | |
| <i>Vidua chalybeata</i> Village Indigobird | | | | | ES | |
| <i>V. macroura</i> Pin-tailed Whydah | C | | | | ES | |
| Fringillidae | | | | | | |
| <i>Serinus mozambicus</i> Yellow-fronted Canary | | | | | <u>EE</u> | |
| Emberizidae | | | | | | |
| <i>Emberiza cabanisi</i> Cabanis's Bunting | | | | | ES | X |
| Totals: 71 families, 398 species (269 in IITA campus IBA; 138 in forest reserve) | 233 | 75 | 74 | | 361 | 25(13) 68(62) |

Discussion

A total of 398 bird species from at least 71 families is now recorded for the Ibadan area, of which the IITA campus IBA holds at least 269 species (68 %) from 64 families, of which 75 species have been mist-netted during our study, while the forest reserve holds at least 138 species, of which 36 species were mist-netted there during our study. Twenty-five species plus 13 vagrants are new to the IITA campus IBA, having not been detected on the site prior to 2002. However, 68 species plus an additional 62 vagrant species reported in the Ibadan area or in IITA by earlier studies have not been detected on the site recently. In general, the diversity of some groups of large birds (*e.g.* Anatidae) has declined, although many of these were vagrants, whereas many forest edge or generalist species (*e.g.* *Francolinus bicalcaratus*) have increased in abundance and range. Forest specialists, including many Pycnonotidae and Bucerotidae, appear to have declined. The use of mist-nets aided the detection of several species previously unknown from the IITA campus, including *Indicator maculatus*, *Campethera nivosa*, *Smithornis rufolateralis*, *Andropadus gracilis*, *Neocossyphus poensis*, *Macrosphenus kempfi*, *M. concolor* and *Cinnyris minullus*.

For the IBA, Ezealor (2001) listed 67 species restricted to the Guinea Congo Rainforest (GCR) biome and seven restricted to the Sudan Guinea Savanna (SGS) biome, with "restricted" meaning having 70 % or more of their range within a particular biome (Fishpool & Evans 2001). We report an additional 29 biome-restricted species of which 27 belong to the GCR (17 of which occur in the forest reserve) and two to the SGS, bringing the total GCR species in the Ibadan area to 94 (84 in the IBA plus 10 without), although nine (plus five classed as vagrants) of the biome-restricted species mentioned by Ezealor (2001) as occurring in the IBA have not been sighted recently (*i.e.* the GCR species *Pteronetta hartlaubii*, *Gymnobucco peli*, *G. calvus*, *Melichneutes robustus*, *Dendropicos gabonensis*, *Coracina azurea*, *Ixonotus guttatus*, *Phyllastrephus icterinus*, *Criniger ndussumensis*, *Cisticola anonymus*, *Muscicapa tessmanni* and *Cinnyris superbus*, and the SGS species *Musophaga violacea* and *Hirundo leucosoma*). These changes, if genuine losses, have occurred over little more than a decade. Ten GCR species, observed in Ibadan by earlier studies but not at the IBA by Ezealor (2001), have not been sighted recently in Ibadan either, *i.e.* *Francolinus lathami*, *Columba iriditorques*, *Prodotiscus insignis*, *Erythrocerus mccallii*, *Megabyas flammulatus*, *Illadopsis puveli*, *Turdoides reinwardtii*, *Antheptes rectirostris*, *Plocepasser superciliosus* and *Nigrita luteifrons*. Nonetheless, the loss of these species depicts the importance of conserving sites such as IITA where their forest habitat is being preserved and now extended. Many of the forest fragments mentioned by Elgood & Sibley (1964) are now a shadow of their former selves, with some now occupied by plantations of fast-growing exotics such as *Gmelina*, *Tectona* and *Eucalyptus* species.

The 17 new GCR species now occurring at the IBA are *Bubo poensis*, *Pogoniulus atroflavus*, *Indicator maculatus*, *Campethera nivosa*, *Smithornis rufolateralis*, *Andropadus*

gracilis, *Cossypha cyanocampter*, *Neocossyphus poensis*, *Macrosphenus kemp*, *M. concolor*, *Sylvietta denti*, *Fraseria ocreata*, *Muscicapa olicascens*, *Cinnyris minullus*, *Oriolus brachyrhynchus*, *Dicrurus atripennis* and *Nigrita fusconotus*. The two new SGS species are *Eremomela pusilla* and *Ptilostomus afer*. The apparent arrival of these species at the site might be attributed to genuine colonisation or to their having been overlooked in previous surveys. The identification of most of these species was confirmed by mist netting, with ringing data available for verification.

The Ibadan area still holds many bird species, although some 68 species (plus 62 vagrants) recorded prior to 2002 have not been recorded recently and now appear to be missing from the area. Reasons may include the fact that some of these are inconspicuous species of high forest, not usually found outside this habitat. However, many others are conspicuous and would therefore not be easily overlooked. A few others may return as vagrants, while a good number are migrants and therefore recorded seasonally.

The comparison of recent with earlier records reveals significant changes in the avifauna of the IITA campus and the surrounding Ibadan area. Some of these changes might not be directly attributed to changes within the campus but rather to the ongoing destruction of forest patches outside it. The city of Ibadan is expanding and many of the patches of forest around the reserve have now been replaced by housing, and natural corridors are thinning out. On the other hand, the extension of the derived savanna up to the edges of the campus IBA, as a result of farmland degradation and clearance of forest and bush, has permitted colonization by an increasing number of birds formerly unknown to the area when it was forested.

Although the IITA campus IBA is protected by a fence from cattle grazers and loggers, the farming activities of the research institute itself are beginning to encroach into its forest reserve areas. Reasons given for this include low productivity from old farm plots, therefore new sites for farm plots are often taken from areas formerly covered by secondary forest. In support of conservation on the site, reforestation over the last four years has begun increasing the area covered by forest though it has not been made clear whether the forest is primarily managed to improve the productivity of soils for agriculture or to conserve soil and wildlife. A balance between land use and biological diversity needs to be agreed upon. Further studies are needed to monitor changes in avifaunal diversity along land use gradients including outside the IITA campus, and longer-term studies are required to monitor bird populations within the IBA.

Acknowledgments

We thank APLORI for collaboration and logistical support during the field work. We acknowledge financial and moral support of the Institute of Immunology, Laboratoire National De Santé, Luxembourg, who provided funding for TAA's PhD program at the University of Ibadan. We thank two anonymous reviewers for their constructive

comments, which greatly improved this report. We thank Matt Stevens, Grace Torkura, Hope Usieta, John Peacock and Ruth Akagu for stimulating companionship and support during field-work. We appreciate John Peacock's enthusiasm for our study, and the hospitality and collaboration of the IITA Management and IITA-Leventis Forest Project to staff and rangers during the field work. This is contribution no. 75 from the A.P. Leventis Ornithological Research Institute.

References

- ADEYANJU T.E & ADEYANJU, T.A (2012) Avifauna of University of Ibadan environs Ibadan, Nigeria. Pp. 27–34 in BABALOLO, F.D., COKER, O.M. & BOROKINI, T.I. (eds) *Proceedings of 3rd Annual Seminar of Nigerian Tropical Biological Association (NTBA)*. Nigerian Tropical Biology Association, Ibadan.
- ASH, J.S. (1990) Additions to the avifauna of Nigeria, with notes on distributional changes and breeding. *Malimbus* 11: 104–116.
- ASHFORD, R.W. (1968) Hirundine records from University of Ibadan campus. *Bull. Niger. Orn. Soc.* 5: 48.
- ASHFORD, R.W. (1969) Two species new to Ibadan and associated records. *Bull. Niger. Orn. Soc.* 6: 30–31.
- ASHFORD, R.W. & PARKER, R.W. (1968) Reed warblers at Ibadan: two species new to the area. *Bull. Niger. Orn. Soc.* 5: 96.
- BANNERMAN, D.A. (1930–51) *The Birds of Tropical West Africa*, vols 1–8. Crown Agents, London.
- BASS, H.G.M. (1967) Notes on birds at Ibadan and elsewhere. *Bull. Niger. Orn. Soc.* 4(13): 31–35.
- BORROW, N. & DEMEY, R. (2010) *Field Guide to the Birds of Western Africa*. Christopher Helm, London.
- BROADBENT, J.A. (1972) Interesting records for the Ibadan area. *Bull. Niger. Orn. Soc.* 8: 54–58.
- BUTTON, J.A. (1965) Harlequin Quail at Illaro and Ibadan. *Bull. Niger. Orn. Soc.* 2: 53.
- DEMEY, R. (2006) Recent Reports. *Bull. Afr. Bird Club* 13: 228.
- DEMEY, R., DOWSETT-LEMAIRE, F. & DOWSETT, R.J. (2003) Notable bird observations from Nigeria, including the first records of Spot-breasted Ibis *Bostrychia rara* and Yellow Longbill *Macrosphenus flavicans*. *Malimbus* 25: 85–94.
- ELGOOD, J.H. & SIBLEY, F.C. (1964) The tropical forest edge avifauna of Ibadan, Nigeria. *Ibis* 106: 221–241.
- ELGOOD, J.H., FRY, C.H. & DOWSETT, R.J. (1973) African migrants in Nigeria. *Ibis* 115: 375–409
- ELGOOD, J.H., HEIGHAM, J.B., MOORE, A.M., NASON, A.M., SHARLAND, R.E. & SKINNER, N.J. (1994) *The Birds of Nigeria*. Checklist 4, 2nd ed., British Ornithologists' Union, Tring.

- EZEALOR, A.U. (2001) Nigeria. Pp. 661–692 in FISHPOOL, L.D.C., & EVANS, M.I. (eds) *Important Bird Areas in Africa and Associated Islands: priority sites for conservation*. Pisces, Newbury.
- FISHPOOL, L.D.C. & EVANS, M.I. (eds) (2001) *Important Bird Areas in Africa and Associated Islands: priority sites for conservation*. Pisces, Newbury.
- MANU, S., PEACH, W. & CRESSWELL, W. (2005) Notes on the natural history of the Ibadan Malimbe *Malimbus ibadanensis*, a threatened Nigerian endemic. *Malimbus* 27: 33–39.
- PARKER, R.H. (1967) Additions to local avifaunas Ibadan. *Bull. Niger. Orn. Soc.* 4(15): 16.
- PARKER, R.H. (1968) Two interesting migrants at Ibadan. *Bull. Niger. Orn. Soc.* 5: 96.
- PARKER, R.H. (1970) Additions to the local avifauna — Ibadan. *Bull. Niger. Orn. Soc.* 7: 38.
- PETTET, A. (1968a) Miscellaneous bird records for the Ibadan area. *Bull. Niger. Orn. Soc.* 5: 52.
- PETTET, A. (1968b) New records for the Ibadan area. *Bull. Niger. Orn. Soc.* 5: 51.
- PETTET, A. (1975) Breeding behavior of *Clamator glandarius* at Ibadan in southern Nigeria. *Bull. Niger. Orn. Soc.* 11: 34–40.
- ROBINSON, D.M. (1966) Wood Ibis at Ibadan. *Bull. Niger. Orn. Soc.* 3: 13.
- ROBINSON, N. (1970) A bird bath in an Ibadan compound: Jericho reservation. *Bull. Niger. Orn. Soc.* 7: 27–30.
- SERLE, W. (1950) Notes on the birds of south-western Nigeria. *Ibis* 92: 84–94.
- THIBAULT, M., DEFOS DU RAU, P. & WELCH, S. (2012) First records of Long-legged Pipit *Anthus pallidiventris* for Nigeria and Benin. *Bull. Afr. Bird Club* 19: 206–208.
- WELLS, D.R. (1966a) A Striped Crake *Porzana marginalis* at Ibadan. *Bull. Niger. Orn. Soc.* 3: 13.
- WELLS, D.R. (1966b) Migrant Scops Owls *Otus scops* at Ibadan. *Bull. Niger. Orn. Soc.* 3: 10–11.
- WELLS, D.R. (1967) Additions to local avifaunas, Ibadan. *Bull. Niger. Orn. Soc.* 4(13/14): 41.
- WILLOUGHBY, R.W.H. (1949) Birds of Ibadan Reservoir. *Niger. Field* 14: 142–145.

L'avifaune d'un milieu de riziculture et de ses environs dans la zone humide de Grand-Bassam, sud-est Côte d'Ivoire

par Guillaume S.K. ODOUKPÉ, Hilaire K. YAOKOKORÉ-BÉIBRO*,
Michael E. KONAN & Pierre K. KOUADIO

Université de Cocody-Abidjan, UFR Biosciences,
Laboratoire de Zoologie et Biologie Animale, BP 582 Abidjan 22, Côte d'Ivoire

*Correspondance: <hyaokokore@yahoo.fr>

Reçu 24 décembre 2013; revu 29 juillet 2014

Résumé

Une étude de l'avifaune a été menée de juillet à septembre 2009 dans le milieu rizicole inondé de la zone humide d'importance internationale de Grand-Bassam. Au cours de ces trois mois, 121 espèces d'oiseaux de 38 familles ont été observées, y compris 30 espèces nouvelles observées pour ce site, ce qui porte la liste de l'avifaune de cette zone humide à 186 espèces.

Summary

Birds of a rice-field area in the Grand Bassam wetlands, south-east Ivory Coast. We undertook a study of the birds in an area of flooded rice cultivation in the Grand-Bassam Wetland of International Importance from July to September 2009. Over the three months, 121 bird species from 38 families were observed, including 30 species new for the site, bringing the avifauna of the wetland to 186 species.

Introduction

Dans un environnement naturel modifié par l'extension des cultures de riz *Oryza* spp., certaines espèces d'oiseaux qui souffraient d'une forte mortalité naturelle en fin de saison sèche trouvent maintenant de quoi survivre et se multiplier, causant d'importants dégâts (Tréca 1985, Manikowski *et al.* 1991). D'autres sont attirées par les cultures et s'y installent pour toute la saison. Les diverses méthodes de protection des cultures ont pour objectif premier de remettre les déprédateurs à leur place. Encore faut-il déterminer les espèces responsables des dégâts, connaître leur comportement et les moments où elles causent le plus de pertes, afin que des aménagements judicieux, l'emploi de bons procédés cultureux et un calendrier

cultural basé sur les dates de présence et d'absence des oiseaux déprédateurs, suffisent à réduire le niveau des dégâts (Tréca 1985, 1989, Manikowski *et al.* 1991).

La culture du riz est pratiquée de plus en plus en Côte d'Ivoire et depuis quelques années la politique gouvernementale en matière d'autosuffisance alimentaire encourage sa production. À Grand-Bassam, certaines espèces d'oiseaux qui fréquentent l'agro-écosystème de riziculture intensive pourraient provoquer des pertes sur leur productivité. Cependant, aucune donnée scientifique sur la thématique n'est disponible aussi bien pour l'avifaune que pour les pertes occasionnées. La base de toute compréhension de la problématique repose sur la connaissance de la diversité des oiseaux qui fréquentent les différents habitats de ce milieu. Le but de cette étude est de connaître la communauté des oiseaux qui fréquentent aussi bien les champs de riz que les divers autres habitats que renferme ce milieu de riziculture.

Site d'étude et méthodes

La zone humide de Grand-Bassam (5°21'N, 3°46'W) est située dans la partie est du littoral ivoirien (Fig. 1). Elle couvre une superficie de 40 210 ha (Yaokokoré-Béibro *et al.* 2010). Le site d'étude, de coordonnées 5°13'N, 3°42'W et 5°14'N, 3°41'W, est une bande de terre comprise entre la lagune Ebrié et le fleuve Comoé. Le climat, de type tropical humide, est caractérisé par deux saisons pluvieuses (avril à mi-juillet et mi-septembre à novembre) et deux saisons sèches (Brou 1997). La pluviométrie varie de 2000 à 2200 mm avec une moyenne annuelle de 2100 mm. La température varie de 21 à 33°C, avec une moyenne annuelle de 26,5°C. La moyenne annuelle de l'humidité relative avoisine 85% (Société d'Exploitation Aérienne et Météorologique: données non-publiées). Le fleuve Comoé et la lagune Ebrié confluent dans cette zone humide pour former le plus vaste estuaire du littoral ivoirien. La région appartient au secteur littoral du grand domaine phytogéographique guinéen et est caractérisée majoritairement par des prairies (75% de la superficie) et des galeries forestières marécageuses, qui se présentent sous la forme de longues bandes étroites parallèles au rivage lagunaire (Guillaumet & Adjanohoun 1971). A coté de ces formations naturelles, le paysage de Grand-Bassam est aussi caractérisé par les plantations de Cocotiers *Cocos nucifera*, de Palmiers à huile *Elaeis guineensis* et de riz (inondé sans régulation de l'eau) *Oryza glaberrima* et *O. sativa*.

En vue d'identifier un grand nombre d'espèces et de couvrir l'ensemble des habitats, le site d'étude a été scindé en cinq zones, selon leur accessibilité. Dans chaque zone, quatre jours d'inventaires ont été effectués mensuellement de juillet à septembre 2009. Ces inventaires ont eu lieu de 6h30 à 11h00 et de 16h00 à 18h00 (Yaokokoré-Béibro 2001). La méthode de points de comptage a permis d'effectuer des inventaires dans les cinq zones.

Pour la reconnaissance des espèces, Borrow & Demey (2001) et les disques des chants et cris de Chappuis (2000) ont été utilisés. L'ordre, la nomenclature et le statut

biogéographique des espèces suivent Borrow & Demey (2001), leurs habitats préférés Yaokokoré-Bébro (2001) et leurs biomes Fishpool & Evans (2001).




Figure 1. Localisation de la zone d'étude indiquant les parcelles rizicoles dans la zone humide de Grand Bassam.

Résultats

Au total, 121 espèces de 37 familles ont été identifiées au cours de l'étude (Tableau 1). Les non-passériformes sont représentés par 63 espèces, et les passériformes par 58 espèces. Le statut biogéographique indique que 99 espèces sont typiquement résidentes. Huit espèces sont typiquement migratrices dont cinq migrateurs paléarctiques, deux migrateurs intra-africains et un migrateur mixte (Tableau 1). Quatorze espèces sont à la fois résidentes et migratrices.

Les espèces inféodées aux milieux ouverts sont les plus nombreuses (58 soit 48 % du peuplement). Elles sont suivies par les espèces forestières généralistes (29 soit 24 % du peuplement) et les espèces des zones humides (23 soit 19 % du peuplement). Les espèces sténotypiques de forêt sont les moins nombreuses (11 soit 9 % du peuplement).

Quarante-sept espèces (soit 39 % des espèces observées) ont été inventoriées aussi bien dans les champs de riz que dans les autres habitats. Les autres 74 espèces (61 %) n'ont pas été inventoriées dans les rizières au cours de l'étude.

Tableau 1. Liste des oiseaux observés dans la zone de riziculture de Grand-Bassam. † nouvellement observée dans la zone. Statut biogéographique: R = résident; P = migrateur paléarctique; M = migrateur intra-africain. Habitat principal: E = eaux, zones humides; O = milieux ouverts; F = forêt secondaire; FF = forêt primaire; * observée dans les champs de riz; A04 = restreinte au biome des savanes soudano-guinéenne; A05 = restreinte au biome des forêts guinéo-congolaises.

| | Statut | Habitat |
|---|--------|---------|
| Phalacrocoracidae | | |
| <i>Phalacrocorax africanus</i> Cormoran africain | R | E |
| Ardeidae | | |
| <i>Ixobrychus minutus</i> Blongios nain | RP | E* |
| <i>Nycticorax nycticorax</i> Bihoreau gris | RP | E* |
| <i>Ardeola ralloides</i> Crabier chevelu | RP | E* |
| <i>Bubulcus ibis</i> Héron garde-bœufs | RM | E* |
| <i>Butorides striata</i> Héron strié | R | E* |
| <i>Egretta gularis</i> Aigrette à gorge blanche | R | E* |
| <i>E. intermedia</i> Aigrette intermédiaire | RM | E* |
| <i>E. alba</i> Grande aigrette | RM | E* |
| Anatidae | | |
| <i>Dendrocygna viduata</i> Dendrocygne veuf* | RM | E* |
| <i>Nettapus auritus</i> Sarcelle à oreillons | R | E* |
| Accipitridae | | |
| † <i>Elanus caeruleus</i> Elanion blanc | R | O* |
| <i>Milvus migrans</i> Milan noir | MP | O |
| <i>Haliaeetus vocifer</i> Pygargue vocifer | R | E |
| <i>Gypohierax angolensis</i> Palmiste africain | R | F |
| <i>Polyboroides typus</i> Gymnogène d'Afrique | R | F |
| <i>Accipiter badius</i> Epervier shikra | RM | O |
| Falconidae | | |
| <i>Falco ardosiaceus</i> Faucon ardoisé | R | O |
| Phasianidae | | |
| <i>Francolinus bicalcaratus</i> Francolin à double éperon | R | O |
| Rallidae | | |
| † <i>Crex egregia</i> Râle des prés | MR | O* |
| <i>Amaurornis flavirostra</i> Râle à bec jaune | R | E* |
| † <i>Porphyrio alleni</i> Talève d'Allen | MR | E* |
| † <i>Gallinula chloropus</i> Gallinule poule d'eau | R | E* |
| Jacanidae | | |
| <i>Actophilornis africanus</i> Jacana à poitrine dorée | R | E* |

| | Statut | Habitat |
|---|--------|---------|
| Rostratulidae | | |
| † <i>Rostratula benghalensis</i> Rhynchée peinte | RM | E* |
| Charadriidae | | |
| <i>Vanellus spinosus</i> Vanneau à éperons | R | E* |
| Scolopacidae | | |
| † <i>Gallinago gallinago</i> Bécassine des marais | P | E* |
| <i>Numenius phaeopus</i> Courlis corlieu | P | E |
| † <i>Tringa glareola</i> Chevalier sylvain | P | E* |
| <i>Actitis hypoleucos</i> Chevalier guignette | P | E |
| Columbidae | | |
| <i>Treron calvus</i> Colombar à front nu | R | F |
| <i>Turtur tympanistria</i> Tourtelette tambourette | R | F |
| <i>T. afer</i> Tourtelette améthystine | R | O* |
| <i>Streptopelia semitorquata</i> Tourterelle à collier | R | O* |
| <i>S. senegalensis</i> Tourterelle maillée | R | O* |
| Psittacidae | | |
| <i>Psittacus erithacus</i> Perroquet jaco | R | FF A05 |
| † <i>Poicephalus senegalus</i> Perroquet youyou | R | O A04 |
| <i>Psittacula krameri</i> Perruche à collier | R | O |
| Musophagidae | | |
| <i>Tauraco macrorhynchus</i> Touraco à gros bec | R | FF A05 |
| <i>Crinifer piscator</i> Touraco gris | R | O |
| Cuculidae | | |
| <i>Chrysococcyx cupreus</i> Coucou foliotocol | R | F |
| <i>C. klaas</i> Coucou de Klaas | R | O |
| <i>C. caprius</i> Coucou didric | R | O* |
| <i>Ceuthmochares aereus</i> Malcoha à bec jaune | R | F |
| † <i>Centropus grillii</i> Coucal de Grill | MR | O |
| <i>C. senegalensis</i> Coucal du Sénégal | R | O |
| Apodidae | | |
| <i>Cypsiurus parvus</i> Martinet des palmiers | R | O |
| <i>Apus affinis</i> Martinet des maisons | R | O |
| Alcedinidae | | |
| † <i>Halcyon malimbica</i> Martin-chasseur à poitrine bleue | R | F |
| <i>H. senegalensis</i> Martin-chasseur du Sénégal | R | O |
| <i>Alcedo cristata</i> Martin-pêcheur huppé | R | O* |
| <i>Ceryle rudis</i> Martin-pêcheur pie | R | E* |
| Meropidae | | |
| <i>Merops pusillus</i> Guépier nain | R | O* |
| Bucerotidae | | |
| <i>Tockus fasciatus</i> Calao longibande | R | F A05 |
| <i>Bycanistes fistulator</i> Calao siffleur | R | FF A05 |
| Capitonidae | | |
| <i>Gymnobucco calvus</i> Barbican chauve | R | F A05 |

| | Statut | Habitat |
|---|--------|---------|
| <i>Pogoniulus scolopaceus</i> Barbion grivelé | R | F A05 |
| † <i>P. atroflavus</i> Barbion à croupion rouge | R | FF A05 |
| <i>P. subsulphureus</i> Barbion à gorge jaune | R | FF A05 |
| <i>P. bilineatus</i> Barbion à croupion jaune | R | F |
| <i>Lybius vieilloti</i> Barbican de Vieillot | R | O |
| † <i>L. bidentatus</i> Barbican bidenté | R | O |
| Picidae | | |
| <i>Dendropicops pyrrhogaster</i> Pic à ventre de feu | R | F A05 |
| Hirundinidae | | |
| <i>Hirundo semirufa</i> Hirondelle à ventre roux | R | O |
| <i>H. abyssinica</i> Hirondelle striée | RM | O |
| <i>H. rustica</i> Hirondelle rustique | P | O |
| Motacillidae | | |
| <i>Anthus leucophrys</i> Pipit à dos uni | R | O* |
| <i>Macronyx croceus</i> Sentinelle à gorge jaune | R | O* |
| Pycnonotidae | | |
| <i>Andropadus virens</i> Bulbul verdâtre | R | F |
| <i>Chlorocichla simplex</i> Bulbul modeste | R | F A05 |
| <i>Thescelocichla leucopleura</i> Bulbul des raphias | R | F A05 |
| <i>Pycnonotus barbatus</i> Bulbul des jardins | R | O |
| Turdidae | | |
| <i>Cossypha niveicapilla</i> Cossyphe à calotte neigeuse | R | F |
| Sylviidae | | |
| † <i>Cisticola erythrops</i> Cisticole à face rousse | R | O* |
| † <i>C. galactotes</i> Cisticole roussâtre | R | O* |
| † <i>C. natalensis</i> Cisticole striée | R | O |
| <i>C. brachypterus</i> Cisticole à ailes courtes | R | O* |
| † <i>Prinia subflava</i> Prinia modeste | R | O* |
| <i>Camaroptera brachyura</i> Camaroptère à tête grise | R | O |
| <i>C. supercilialis</i> Camaroptère à sourcils jaunes | R | FF A05 |
| † <i>C. chloronota</i> Camaroptère à dos vert | R | FF A05 |
| † <i>Sylvietta virens</i> Fauvette Crombec verte | R | F A05 |
| <i>Hylia prasina</i> Hylia verte | R | F A05 |
| Monarchidae | | |
| <i>Terpsiphone rufiventer</i> Tchitrec à ventre roux | R | F A05 |
| Platysteiridae | | |
| <i>Platysteira castanea</i> Pririt châtain | R | FF A05 |
| <i>P. cyanea</i> Pririt à collier | R | O |
| Remizidae | | |
| † <i>Pholidornis rushiae</i> Méсанgette rayée | R | O A05 |
| Nectariniidae | | |
| † <i>Anthreptes rectirostris</i> Souimanga à bec droit | R | FF A05 |
| <i>Anabathmis reichenbachii</i> Souimanga de Reichenbach | R | O A05 |
| <i>Cyanomitra verticalis</i> Souimanga olive à tête bleue | R | F |

| | Statut | Habitat |
|---|--------|---------|
| † <i>Chalcomitra adelberti</i> Souimanga à gorge rousse | R | FF A05 |
| <i>C. senegalensis</i> Souimanga à poitrine rouge | RM | O |
| <i>Hedydipna collaris</i> Souimanga à collier | R | F |
| <i>Cinnyris chloropygius</i> Souimanga à ventre olive | R | F |
| † <i>C. coccinigastrus</i> Souimanga éclatant | R | O A04 |
| <i>C. cupreus</i> Souimanga cuivré | R | O* |
| Laniidae | | |
| <i>Lanius collaris</i> Pie-grièche fiscale | R | O* |
| Malaconotidae | | |
| † <i>Dryoscopus gambensis</i> Cubla de Gambie | R | F |
| Corvidae | | |
| <i>Corvus albus</i> Corbeau pie | R | O |
| Sturnidae | | |
| <i>Lamprotornis splendidus</i> Choucador splendide | R | F |
| † <i>Cinnyricinclus leucogaster</i> Spréo améthyste | M | O |
| Passeridae | | |
| <i>Passer griseus</i> Moineau gris | R | O* |
| Ploceidae | | |
| † <i>Ploceus pelzelni</i> Tisserin de Pelzeln | R | O |
| <i>P. nigricollis</i> Tisserin à cou noir | R | O |
| <i>P. aurentius</i> Tisserin orangé | R | O A05 |
| <i>P. nigerrimus</i> Tisserin noir de Vieillot | R | O* A05 |
| <i>P. cucullatus</i> Tisserin gendarme | R | O* |
| <i>Malimbus nitens</i> Malimbe à bec bleu | R | F A05 |
| <i>M. scutatus</i> Malimbe à queue rouge | R | FF A05 |
| † <i>Quelea erythrops</i> Travailleur à tête rouge | M | O* |
| † <i>Euplectes afer</i> Euplecte vorabé | R | O* |
| <i>E. macrourus</i> Euplecte à dos d'or | R | O* |
| Estrildidae | | |
| <i>Nigrita canicapillus</i> Nigrette à calotte grise | R | F |
| † <i>N. bicolor</i> Nigrette à ventre roux | R | FF A05 |
| <i>Pyrenestes ostrinus</i> Pyréneste ponceau | R | F* |
| † <i>Spermophaga haematina</i> Sénégal sanguin | R | F A05 |
| <i>Estrilda melpoda</i> Astrild à joues orange | R | O* |
| <i>Lonchura cucullata</i> Capucin nonnette | R | O* |
| <i>L. bicolor</i> Capucin bicolore | R | O* |
| † <i>L. fringilloides</i> Capucin pie | R | O* |
| Viduidae | | |
| <i>Vidua macroura</i> Veuve dominicaine | R | O* |

Quatre espèces méritent des commentaires compte tenu du peu de données dont on dispose sur elles dans cette région. Un individu de *Centropus grillii* a été observé le

10 août 2009 dans la zone 5. La présence de l'espèce est confirmée par les travaux de Gueye (2013). Cette espèce se distingue de la forme *epomidis* de *C. senegalensis*, aussi présente et résidente dans la région, par sa coloration noire (tête, parties inférieures, queue) et ses ailes roussâtres. La forme *epomidis*, assez commune entre les individus typiques de *C. senegalensis* sur le site et bien connu des auteurs, se caractérise par la tête et la gorge noires et par les ailes et les parties inférieures marron. Deux individus de *Lybius bidentatus* ont été régulièrement observés dans les zones 1 et 3, de juillet à septembre 2009. *Lybius bidentatus* est semblable à *L. dubius*, mais s'en distingue par l'absence de bande pectorale noire et la présence d'une barre alaire rouge. Cette espèce a déjà été mentionnée dans le sud-est de la Côte d'Ivoire, dans le parc national des îles Ehotilé (Yaokokoré-Béibro 2010). Un individu de *Chalcomitra senegalensis* a été observé le 19 juillet 2009 dans la zone 1. Cette espèce a également été mentionnée dans cette région par Yaokokoré-Béibro *et al.* (2010). *Pyrenestes ostrinus* a été régulièrement observé dans le site d'étude de juillet à septembre 2009, avec une abondance avoisinant la cinquantaine d'individus. Les mâles adultes de *P. ostrinus* sont semblable à ceux de *P. sanguineus*, que nous n'avons pas enregistré sur le site, par la tête, la poitrine, les flancs, le croupion et la queue rouges ainsi qu'un cercle oculaire blanc; ils sont cependant distincts par le reste de leur coloration qui est noire pour *P. ostrinus* et brun terre pour *P. sanguineus*.

Discussion

La richesse spécifique de la zone prospectée montre que ce milieu de riziculture renferme une grande diversité d'oiseaux. Les travaux antérieurs (Demey & Fishpool, 1991, N'guessan 2007, Lachenaud *et al.* 2008, Yaokokoré-Béibro *et al.* 2010) ont permis d'inventorier 156 espèces pour la zone humide d'importance internationale de Grand Bassam. L'étude a permis d'identifier 30 espèces nouvelles pour la zone. La liste des espèces de la zone humide de Grand-Bassam s'élève ainsi à 186 espèces dont deux, *Sterna balaenarum* et *Bycanistes cylindricus*, sont proches de la menace (NT) dans la liste rouge de l'UICN (BirdLife International 2004). Toutefois, 45 espèces mentionnées dans la région d'étude (Yaokokoré *et al.* 2010), comprenant de nombreux migrateurs, n'ont pas été observées. Ceci serait lié à la période de l'étude qui est très courte (de juillet à septembre 2009) et défavorable à l'observation des migrateurs paléarctiques (seulement huit migrateurs paléarctiques sont présentés dans cette étude). Vu qu'aucune véritable étude de population de l'avifaune n'a été effectuée dans la région, il est probable que le statut de certaines espèces d'oiseau, telles que *Egretta alba* et *Centropus grillii*, ne soit pas clairement défini. Certaines autres, indiquées comme à la fois résidentes et migratrices, pourraient être seulement résidentes dans la région d'étude, et migratrices dans d'autres parties du pays (*Accipiter badius*, *Rostratula benghalensis*, *Hirundo abyssinica*). Pour d'autres, l'existence d'une population résidente est prouvée, mais on soupçonne également celle de migrateurs

africains (*Egretta intermedia*), ou paléarctiques (*Ixobrychus minutus*, *Nycticorax nycticorax*) ou les deux (*Ardeola ralloides*). *Milvus migrans* est un migrateur africain nicheur dans la région, avec la présence possible de migrateurs paléarctiques.

Trois des espèces typiques qui fréquentent la végétation côtière et les mangroves (*Anabathmis reichenbachii*, *Ploceus aurantius* et *P. pelzelni*) ont été observées, 27 des 188 espèces restreintes au biome des forêts guinéo-congolaises et deux des 38 espèces restreintes au biome des savanes soudano-guinéenne (Fishpool & Evans 2001).

Comparé aux autres sites du littoral ivoirien, la liste totale des 186 espèces d'oiseaux du milieu rizicole de Grand Bassam partage 83 % de son peuplement avec la forêt classée de N'ganda N'ganda (Kouadio 2006), 72 % avec le parc national des Iles Ehotilé (Yaokokoré-Béibro 2010) et 55 % avec le parc national d'Azagny (Fishpool & Evans 2001, Demey 2006). La zone humide de Sassandra-Dagbégo, plus au sud-ouest, partage avec le site d'étude 59 % du peuplement (Lachenaud 2006). Ces chiffres montrent que la zone humide Ramsar de Grand-Bassam est bien complémentaire des autres sites du littoral ivoirien et vaut d'être protégée.

Cette liste préliminaire du peuplement d'oiseaux dans la zone de riziculture de la zone humide de Grand-Bassam montre que les espèces qui fréquentent les champs cultivés ne sont pas toutes des déprédatrices du riz. Certaines sont insectivores et contribuent à la régulation naturelle des insectes ravageurs des cultures. D'autres sont des prédateurs d'oiseaux, de rongeurs et de reptiles. La lutte ou la protection des champs contre les oiseaux nécessite donc de connaître les espèces responsables des pertes et leurs caractéristiques écologiques, biologiques et comportementales. Des études complémentaires sur deux cycles annuels sont nécessaires pour compléter ces premières données.

Bibliographie

- BIRDLIFE INTERNATIONAL. (2004) *Threatened Birds of the World 2004*. CD-ROM, BirdLife International, Cambridge.
- BORROW, N. & DEMEY, R. (2001) *Birds of Western Africa*. Christopher Helm, Londres.
- BROU, Y. (1997). Analyse et dynamique de la pluviométrie en milieu forestier ivoirien. Thèse de Doctorat 3ème cycle. Université de Cocody-Abidjan, Abidjan.
- CHAPPUIS, C. (2000) *African Bird Sounds. Birds of North, West and Central Africa*. Livrette et 15 CD. Société d'Etudes Ornithologiques de France, Paris.
- DEMEY R. & FISHPOOL L.D.C. (1991) Additions and annotations to the avifauna of Côte d'Ivoire. *Malimbus*, 12: 61–86.
- DEMEY, R. (2006) Liste des espèces d'oiseaux des parcs nationaux de Côte-d'Ivoire. Pp. 517–532 in LAUGNIE, F. (ed.) 2007. *Conservation de la Nature et Aires Protégées en Côte-d'Ivoire*. NEI/Hachette et Afrique Nature, Abidjan.
- FISHPOOL, L.D.C. & EVANS, M.I. (eds) (2001) *Important Bird Areas in Africa and Associated Islands*. BirdLife International, Cambridge.

- GUEYE M.F. (2013) *Inventaire Ornithologique d'un Milieu Urbain Côtier de Côte d'Ivoire: cas de la ville de Grand-Bassam*. DEA Écologie Tropicale, Université Félix Houphouët-Boigny, Abidjan.
- GUILLAUMET, J.-L. & ADJANOHOON, E. (1971) La végétation de la Côte d'Ivoire. *Mém. Off. Rech. Sci. Tech. Outre-Mer* 50: 157–263.
- KOUADIO, K.P. (2006) Avifaune de la forêt classée de N'ganda N'ganda à Assinie-mafia (Côte d'Ivoire): inventaire et distribution. DEA Écologie Tropicale, Université de Cocody-Abidjan, Abidjan.
- LACHENAUD, O. (2006) Les oiseaux de la région de Sassandra, Côte d'Ivoire. *Malimbus* 28: 18–34.
- LACHENAUD, O., OOSTERHUIS, H.-J., BOEDTS, B. & GUNNINGHAM, J. (2008) Nouvelles observations ornithologiques en Côte d'Ivoire. *Malimbus* 30: 19–29.
- MANIKOWSKI, S., NDIAYE, A.B. & TRÉCA, B. (1991) *Manuel de Protection des Cultures contre les Dégâts d'Oiseaux*. Rapport FAO–Appui à la lutte antiaviaire - Projet TCP/SEN/0053, FAO, Rome.
- N'GUESSAN, A.M. (2007) *Avifaune de la Zone Humide d'Importance Internationale de Grand-Bassam (Côte d'Ivoire): Inventaire, Structure du Peuplement et Origine Biogéographique*. DEA Écologie Tropicale, Université de Cocody-Abidjan, Abidjan.
- THIOLLAY, J.M. (1986) Structure comparée du peuplement avien des trois sites de forêt primaire en Guyane. *Terre Vie* 41: 59–105.
- TRÉCA, B. (1985) Les possibilités de lutte contre les oiseaux d'eau pour protéger les rizières en Afrique de l'Ouest. *J. Agric. trad. Bot. appl.* 23: 191–213.
- TRÉCA, B. (1989) Les risques de dégâts d'oiseaux sur les rizières sahéliennes. Pp. 167–175 in ELDIN, M. & MILLEVILLE, P. (eds), *Le Risque en Agriculture*. ORSTOM, Paris.
- TRÉCA, B. (1992) Waterbirds and rice cultivation in West Africa. *Proc. VII Pan-Afr. Orn. Congr.*: 297–301.
- YAO KOKORÉ-BÉIBRO, K.H. (2001) *Avifaune des Forêts Classées de l'Est de la Côte d'Ivoire: Données sur l'Ecologie des Espèces et Effet de la Déforestation sur les Peuplements. Cas des forêts classées de la Béki et de la Bossématié (Abengourou)*. Thèse de Doctorat, Université de Cocody-Abidjan, Abidjan.
- YAO KOKORÉ-BÉIBRO, K.H. (2010) Oiseaux du Parc National des Iles Ehotilé, sud-est Côte d'Ivoire. *Malimbus* 32: 89–102.
- YAO KOKORÉ-BÉIBRO, K.H., N'GUESSAN, A.M., ODOUKPÉ, K.S.G., ZOZOU, E.J., N'DOUBA, V. & KOUASSI, K.P. (2010) Premières données sur les oiseaux de la zone humide d'importance internationale de Grand-Bassam (Côte d'Ivoire). *Int. J. biol. chem. Sci.* 4: 2169–2180.

Short Notes — Notes Courtes

Records of African Pygmy Goose *Nettapus auritus* and a *Halcyon* kingfisher on São Tomé

On 21 July 2011, one of us (AT) observed a female African Pygmy Goose *Nettapus auritus* feeding and resting in a sheltered part of the river Io Grande, Dona Eugénia area (0.13322°N, 6.61979°E), in southeast São Tomé. The bird was observed for at least 20 min. (10h10–10h30) and several photos were taken (Fig. 1). The distinct plumage of the species and the perfect observational circumstances exclude misidentification. The African Pygmy Goose is distributed over large parts of sub-Saharan Africa, where it is largely confined to freshwater lakes, marshes and rivers (Brown *et al.* 1982, Borrow & Demey 2001). It is noted as vagrant to the islands of São Tomé and Príncipe in the Gulf of Guinea on the IUCN Red List (<<http://www.iucnredlist.org/>>, consulted Feb 2014), but to the best of our knowledge no observations have been formally reported to date. São Tomé is located *c.* 175 km west of the coast of Gabon, and the present record thus demonstrates the species' capability of dispersal over the sea.

The day before this observation, *i.e.* 20 July 2011, we encountered yet another unexpected bird for São Tomé: a *Halcyon* kingfisher. This observation was made during a walk along the Io Grande, close to its junction with the river Ana Chaves (0.13738°N, 6.61971°E). The bird perched on a dead branch of a tree located up the almost vertical bank of the river. Unfortunately, it was raining at the time of the observation, which, in combination with the long distance (*c.* 150 m), rendered detailed observations impossible. We observed clearly the typical silhouette of a *Halcyon* kingfisher, with proportionally very long bill and short tail; the belly and underparts appeared light and the back and head dark; a light patch was observed on the right wing of the bird. The Blue-breasted Kingfisher *H. malimbica* is resident on the island of Príncipe, *c.* 145 km northeast of São Tomé) as a local subspecies *H. m. dryas*, which is slightly larger than its mainland counterpart. It occurs in a wider range of habitats on Príncipe than on the mainland (Christy & Clark 1998). Our observation could possibly have been of this subspecies, but we can neither exclude the mainland subspecies nor other *Halcyon* species (in particular the Woodland Kingfisher *H. senegalensis*). Jones & Tye (2006) considered all previous records of *H. m. dryas* for São Tomé as subject to doubt and concluded that there was no good evidence that the species had truly occurred there; although our observation does not change this situation, it indicates a need for vigilant attention to large kingfishers seen on São Tomé.

We are grateful to the National Geographic Society for funding (Waitt Grant W172-11) and Martim Melo, Octávio Veiga and Yelli for field work.



Figure 1. Female African Pygmy Goose in a sheltered part of the river Io Grande, southeast São Tomé, 21 July 2011 (photo: A. Thomasson).

References

- BORROW, N. & DEMEY, R. (2001) *Birds of Western Africa*. Christopher Helm, London.
- BROWN, L.H., URBAN, E.K. & NEWMAN, K. (1982) *The Birds of Africa*, vol. 1. Academic Press, London.
- CHRISTY, P. & CLARKE, W.V. (1998) *Guide des Oiseaux de São Tomé et Príncipe*. ECOFAC, São Tomé.
- JONES, P. & TYE, A. (2006) *The Birds of São Tomé and Príncipe with Annobón, Islands of the Gulf of Guinea*. British Ornithologists' Union, Oxford.

Received 12 September 2013

Revised 16 February 2014

Bengt HANSSON¹ & August THOMASSON²

¹Department of Biology, Lund University, SE-223 62 Lund, Sweden
<bengt.hansson@biol.lu.se>

²Sösdala 2094, 28010 Sösdala, Sweden

Observations ornithologiques au Sénégal et en Gambie

Pendant un voyage d'ornithologues amateurs parrainé par l'Association des Naturalistes des Yvelines (ANY, France) au Sénégal et en Gambie du 2 au 19 février 2014, les participants ont fait cinq observations particulières.

***Ciconia ciconia* Cigogne blanche.** Nous avons observé environ 1500 Cigognes blanches dans quatre groupes dans la Réserve de Baobolong (13°34'N, 15°47'W) en

Gambie, le 16 fév; des regroupements de centaines ou milliers d'individus de cette espèce ont été signalés pendant les migrations (Géroudet 1994).

***Gorsachius leuconotus* Bihoreau à dos blanc.** Nous avons aussi observé dans la Réserve de Baobolong (13°28'N, 15°48'W), le 14 fév, un adulte et un juvénile du Bihoreau à dos blanc (Fig. 1A). Bien que cette espèce soit déjà connue de cette région, il n'existe que peu d'évidence de sa reproduction.

***Ardea cinerea* Héron cendré.** Un individu de la sous-espèce *monicae* (Héron pâle) observé en compagnie d'autres Ardeidae, dans les arbres bordant la mare de Leba (13°21'N, 13°22'W), près de Badi à l'extérieur du Parc National du Niokolo-Koba, le 11 fév. Les premiers contacts visuels faisaient penser à un Héron cendré leucistique, mais les détails obtenus grâce au télescope ont permis de bien remarquer les taches noires le long du cou, comme illustré dans Borrow & Demey (2011). Cette sous-espèce, considérée par certains comme espèce distincte, ne se trouve normalement qu'aux alentours du Banc d'Arguin en Mauritanie (Borrow & Demey 2011).

***Acrocephalus paludicola* Phragmite aquatique.** Un individu observé sur la mare de Leba, le 11 fév. Deux caractéristiques rendent crédible l'observation de ce phragmite: le dessin bien visible de la tête, avec ses trois bandes claires, nettement détachées, et le milieu dans lequel l'oiseau a été observé, une mare entourée de plantes marécageuses. Lors de l'observation, le phragmite était perché momentanément sur un roseau. Jusqu'ici ce migrateur paléarctique n'a été observé que sur la côte à l'extrême nord du Sénégal (Le Nevé *et al.* 2013).

***Prionops plumatus* Bagadai casqué.** Un groupe d'une dizaine suivaient de très près notre groupe lors d'une promenade à pied dans la bananeraie (13°21'N, 13°21'W) de Wassadou proche du Parc National du Niokolo-Koba, le 3 fév. Les oiseaux étaient agités, volant bas et se posant souvent à quelques mètres des participants (Fig. 1B) tout au long d'un parcours de plus d'une centaine de mètres. Typiquement les Bagadai



Figure 1. A: Bihoreau à dos blanc *Gorsachius leuconotus* (juvénile), 14 fév 2014. **B:** *Prionops plumatus* Bagadai casqué, 3 fév 2014. (Clichés: J. Rose).

casqués fuient quand ils sont approchés de moins d'une vingtaine de mètres, et les guides du Parc national nous ont dit qu'ils n'avaient jamais auparavant vu ce comportement. Peut-être les oiseaux suivaient-ils un chemin habituel.

Nous remercions Lamin Sanyang, conservateur du Parc National de Niomi, qui nous a accompagnés lors de notre visite en Gambie, et Jean-Louis Faure, membre du Groupe Ornithologique Normand, pour sa contribution à la validation des observations.

Bibliographie

- BORROW, N. & DEMEY, R. (2011) *Birds of Senegal and The Gambia*. Christopher Helm, Londres.
- GÉROUDET, P. (1994), *Grands Échassiers, Gallinacées, Râles d'Europe*. Delachaux et Niestlé, Lausanne.
- LE NEVÉ, A., BARGAIN, B., TEGETMEYER, C. & GUYOT, G. (2013) Hivernage au Sénégal d'un Phragmite aquatique *Acrocephalus paludicola* "isabelle": longévité de l'espèce et liens de migration. *Malimbus* 35: 142–146.

Reçu 31 mai 2014

Revu 11 août 2014

Moussa Séga DIOP¹, Bertrand MENNESSON² & John ROSE³

¹Cité SICA de Mbao Villa N° 191, BP 20077 Dakar-Thiaroye, Sénégal;
<msemiop@gmail.com>

²Co-coordonateur d'ornithologie, ANY,

Villa de Chèvreloup, 34 route de Versailles, 78150 Rocquencourt, France

³1 Bis rue des Châtre-Sacs, 92310 Sèvres, France

The Parelus bird collection from Ivory Coast at the Field Museum of Natural History, and the first country record of Rufous Cisticola *Cisticola rufus*

Museum collections can play a critical role in unravelling a country's faunistic history. Each specimen creates a scientific, historical record of where and when a bird occurred, from which, among other things, we can learn about changes in land cover and climate, and infer changes in an organism's distribution over time (e.g. Moritz *et al.* 2008). This is especially useful for difficult to identify species.

We report on a collection of birds from Ivory Coast made by D. Parelus between 1964 and 1969, which has been overlooked in previous treatments of the country's avifauna (Thiollay 1985, Dowsett 1993, Dowsett *et al.* 2014). Parelus, the son of American missionaries, lived in Ivory Coast intermittently during this period. After being given a book on preparing bird specimens by Jim Gould, another missionary in

the area, and corresponding with Melvin Traylor, curator of birds at the Field Museum of Natural History (FMNH: Chicago, U.S.A.), he started collecting birds for the museum in 1964. The first part of the collection (339 specimens collected in 1964–5), published by Traylor & Parelius (1967), added 44 species to the country's known avifauna. In total, the Parelius collection comprises 1933 specimens of 297 species, predominantly from the areas around Bouaké, Abidjan and Korhogo. However, while Parelius himself is acknowledged in the first checklist of birds of Ivory Coast (Brunel & Thiollay 1969), his collection was not examined by those authors (J.-M. Thiollay pers. comm.).

We examined the complete list of Parelius specimens as catalogued and compared it to the most recent checklist of the birds of Ivory Coast (Dowsett *et al.* 2014). After accounting for taxonomic discrepancies, we then examined the specimens of the three species in the catalogue that were not included in Dowsett *et al.* (2014): Rufous Cisticola *Cisticola rufus*, Black-bellied Seedcracker *Pyrenestes ostrinus* and Golden-breasted Bunting *Emberiza flaviventris*. We determined that the bunting specimen was actually Cabanis's Bunting *E. cabinisi*, a species already known from Ivory Coast. However, the other two were correctly identified and we include details of those records here.

***Cisticola rufus* Rufous Cisticola.** These specimens represent the only records of the species for Ivory Coast, heretofore unpublished. Distinguishing Rufous Cisticola from the sympatric Short-winged Cisticola *C. brachypterus* is difficult, obscured by variation in the colour of the upperparts. In addition, based on examination of the Parelius specimens and others in the Field Museum's collections, we found the shape of the



Figure 1. Small cisticolas collected in northern Ivory Coast by D. Parelius. The six on the left are *Cisticola rufus*, the only specimens of that species from the country. The five on the right are *C. brachypterus*.

Table 1. FMNH accession numbers and data for the specimens of *Cisticola rufus* collected by D. Parelus in Ivory Coast. The coordinates are modified from the original field notes and from Traylor & Parelus (1969).

| Accession number | Locality | Coordinates | Date | Sex |
|------------------|-----------|-----------------------|-------------|--------|
| 278904 | Ferké | 9°36'N, 5°12'W | 11 Aug 1966 | Female |
| 278906 | Ferké | 9°36'N, 5°12'W | 5 Aug 1966 | Male |
| 278907 | Boundiali | 9°32'N, 6°29'W | 16 Jul 1966 | Male |
| 278908 | Ferké | 9°36'N, 5°12'W | 6 Aug 1966 | Female |
| 285844 | Boundiali | 9°32'N, 6°29'W | 12 Aug 1966 | Female |
| 285847 | Torgokaha | 1 km west of Korhogo* | 13 Jul 1969 | Male |

*Coordinates for Korhogo: 9°27'N, 5°39'W

outermost primary, considered diagnostic by Lynes (1930), to be unreliable as an identification criterion. However, Rufous *Cisticola* is distinguishable by its more rufous upperparts, apparent in side-by-side comparisons, and by the lack of any hint of streaking on the back (Fig. 1). Based on these features, we have identified six specimens, which Parelus collected from three localities in northern Ivory Coast, as Rufous *Cisticola* (Table 1).

***Pyrenestes ostrinus* Black-bellied Seedcracker.** Four specimens of this widespread estrildid were collected at Port Bouet in March 1967. This taxon is not listed in any checklist of birds of Ivory Coast (e.g. Thiollay 1985, Dowsett 1993, Dowsett *et al.* 2014), perhaps stemming from Brunel & Thiollay's (1969) listing only "*Pyrenestes (ostrinus) sanguineus*", implying that only Crimson Seedcracker *P. sanguineus* was known from the country. This was in spite of an earlier publication by Brunel (1955) stating that he collected nests of both seedcracker species at Bingerville. Regardless, there are several recently published records from the southeast of the country: Grand Bassam (Borrow 2000, Yaokokoré-Béibro *et al.* 2010), Parc National des Iles Ehotilé (Yaokokoré-Béibro 2010) and N'Ganda N'Ganda (Kouadio *et al.* 2014). The Parelus skins may represent the only specimens of this taxon for Ivory Coast.

Not incorporating such distributional data in country checklists can have a residual effect in the literature. For example, Salewski *et al.* (2001) used the range descriptions of Thiollay (1985) as a baseline for detecting latitudinal range shifts in Ivorian birds. However, two of the species listed as potentially having northward range expansions (Blackcap Babbler *Turdoides reinwardtii* and Black-necked Weaver *Ploceus nigricollis*) would have been omitted had Thiollay (1985) incorporated the Parelus collection in his paper. In this case, this is not a serious error: indeed it would have served to strengthen the conclusion of Salewski *et al.* (2001).

This paper is not an exhaustive review of the Parelus collection. Its data are freely available in online museum collection databases (e.g. VertNet, ORNIS). We encourage researchers interested in bird distributions in the Ivory Coast and West Africa in general to incorporate this valuable collection into future studies.

This work was possible thanks to grants from the National Science Foundation for databasing the FMNH bird collection.

References

- BORROW, N. (2000) Recent Reports. *Bull. Afr. Bird Club* 7: 147.
- BRUNEL, J. (1955) Observations sur les oiseaux de la Basse Côte d'Ivoire. *Oiseau Rev. fr. Orn.* 25: 1–16.
- BRUNEL, J. & THIOLLAY, J.-M. (1969) Liste préliminaire des oiseaux de Côte d'Ivoire. *Alauda* 37: 230–254, 315–337.
- DOWSETT, R.J. Ivory Coast. In DOWSETT, R.J. & DOWSETT-LEMAIRE, F. (1993) A contribution to the distribution and taxonomy of Afrotropical and Malagasy birds. *Tauraco Res. Rep* 5: 43–49.
- DOWSETT, R.J., ATKINSON, P.W. & CADDICK, J.A. (2014) Checklist of the birds of Côte d'Ivoire. <www.africanbirdclub.org/countries/checklists/downloads>, consulted 27 March 2014.
- KOUADIO, K.P., YAOKOKORÉ-BÉIBRO, K.H., SAINT GUILLAUME, K., ODOUKPÉ, E.M.K., N'GNUSSAN, A.M. & KOUASSI, K.P. (2014). Diversité avifaunique de la forêt classée de N'ganda N'ganda (sud-est de la Côte d'Ivoire). *Afr. Sci.* 10: 181–193.
- LYNES, H. (1930) Review of the genus *Cisticola*. *Ibis* 6(Suppl.): 1–673.
- MORITZ, C., PATTON, J.L., CONROY, C.J., PARRA, J.L., WHITE, G.B. & BEISSINGER, S.R. (2008) Impact of a century of climate change on small-mammal communities in Yosemite National Park, USA. *Science* 322: 261–264.
- SALEWSKI, V., RAINEY, H. & BAIRLEIN, F. (2001) Have birds shifted their range limits southwards in Côte d'Ivoire, West Africa? *Bull. Afr. Bird Club* 8: 117–120.
- THIOLLAY, J.-M. (1985) The birds of Ivory Coast: status and distribution. *Malimbus* 7: 1–59.
- TRAYLOR, M. & PARELIUS, D. (1967) A collection of birds from the Ivory Coast. *Fieldiana* 51: 91–117.
- YAOKOKORÉ-BÉIBRO, H.K. (2010) Oiseaux du Parc National des Iles Ehotilé, sud-est Côte d'Ivoire. *Malimbus* 32: 89–102.
- YAOKOKORÉ-BÉIBRO, H.K., N'GNUSSAN, M.A., ODOUKPÉ, G.K., ZOUZOU, E.J., N'DOUBA, V. & KOUASSI, P.K. (2010) Premières données sur les oiseaux de la zone humide d'importance internationale de Grand-Bassam (Côte d'Ivoire). *Internat. J. biol. chem. Sci.* 4: 2169–2180.

Received 3 May 2014

Revised 4 July 2014

Joshua I. ENGEL^{1,2}, Daniel PARELIUS³ & John M. BATES¹

¹Field Museum of Natural History, Integrative Research Center,
1400 South Lake Shore Drive, Chicago, Illinois 60605, U.S.A.

²Correspondence <jjengel2@fieldmuseum.org>

³427 Bakers Branch Road, Louisa, Virginia 23093, U.S.A.

News & Comment — Nouvelles & Commentaires

Ornithological services of the Niokolo-Koba National Park guides

Niokolo-Koba National Park (NKNP) in southeast Senegal is, with an area of 913,000 ha, the largest nature sanctuary in that country and one of the largest and most important in West Africa. The exceptional diversity of the fauna and flora of the Park were recognized in 1981 with its designation by UNESCO as a Biosphere Reserve and World Heritage site.

NKNP is also a key resource for local sustainable development and environmental action in the area. With the encouragement of the national and local authorities, the 31 eco-guides of the Park have been working tirelessly through their cooperative called GIE NIOKOLO <<http://www.niokolo-safari.com>> to improve and promote the Park and foster development of the region.

The wealth of bird life in and around NKNP is exceptional, estimated at about 345 species from the distribution maps of Borrow & Demey (2011, *Birds of Senegal and The Gambia*, Christopher Helm, London), without counting isolated or doubtful observations. Since 2012 the French association “COMETE International” <<http://www.comete-international.org>> has been assisting GIE NIOKOLO to improve its tourist services and to contribute more effectively to the advancement and dissemination of scientific knowledge. The guides have thus been able to organise visits to the Park and its surroundings for international groups of amateur ornithologists in 2013 and 2014 (Diop *et al.* 2014, *Malimbus* 36: 117–119; <<http://www.siteany78.org/>>), and plan a third in Senegal and Gambia in 2015. GIE NIOKOLO also offers birdwatching treks in areas of interest outside the Park, notably just north of it (near the main entrance) and in the foothills near the Guinean border to the south. Starting in November 2014, the GIE will be systematically collecting and posting data from its ornithological observations on its website, with an emphasis on raptors, water birds and species of particular interest for ornithological tourism.

To participate in sharing information on the birds of NKNP, or for questions concerning the ornithological services of GIE NIOKOLO, readers are invited to contact us.

Ibrahima KOUYATÉ¹, Moussa Séga DIOP² & John ROSE³

¹Chair of the Ornithology Commission of GIE NIOKOLO, BP 362, Tambacounda, Sénégal. <info@niokolo-safari.com>

²Trustee of COMETE International, Cité SICA de Mbao Villa N° 191, BP 20077 Dakar-Thiaroye, Sénégal. <mstediop@gmail.com>

³President of COMETE International, Issy-les-Moulineaux, France. <comete@comete-international.org>

Society Notices — Informations de la Société

W.A.O.S. membership changes Changements à la liste d'adhérents de la S.O.O.A.

New members — Nouveaux membres

FOX, O., Clock House, Coleshill, Swindon, Wiltshire SN6 7PT, **U.K.**

GUÉVORTS, B., Rue du Prince de Galles, Akwa, BP 15779 Douala, **Cameroon**

MUNDY, R., 9 Market Lane, Lewes, East Sussex BN7 2DS, **U.K.**

OSINUBI, S.T., Swansley Wood Farm, Cambridge Rd., Caxton, Cambridge CB23 3PH,
U.K.

PARELIUS, D., 427 Bakers Branch Rd, Louisa, VA 23093, **U.S.A.**

NATURHISTORISCHES MUSEUM, Burgring 7, A-1010 Vienna, **Austria**

Address changes and corrections — Changements et corrections d'adresse

COWAN, Dr P., 3 Bank Manor, Bank Road, Matlock DE4 3NF, **U.K.**

FRY, Prof. C.H., Bridge End House, Kentmere, Kendal, Cumbria LA8 9JP, **U.K.**

KING, T., Cantuar, High Street, Lyminge, Kent CT18 8EN, **U.K.**

KOUGOUM, P.G., Dept of Biological Sciences, Faculty of Sciences, University of
Maroua, PO Box 814, Maroua, **Cameroon**

WILKINSON, Dr R., 2 Weston Grove, Upton-by-Chester, Chester, Cheshire CH2 1QJ,
U.K.

WILSON ORNITHOLOGICAL SOCIETY, Museum of Zoology, The University of
Michigan, 1109 Geddes Ave, Ann Arbor, MI 48109-1079, **U.S.A.**

Tim Dodman, Treasurer and Membership Secretary

**West African Ornithological Society
Société d'Ornithologie de l'Ouest Africain**



Membership List, 1 February 2014

West African Ornithological Society Soci t  d'Ornithologie de l'Ouest Africain

Membership List, 1 February 2014

Would members please check their entries and notify any mistakes or changes to the Treasurer and Membership Secretary: Tim Dodman, Hundland, Papa Westray, Orkney KW17 2BU, U.K. <tim@timdodman.co.uk>.

To facilitate communication with members and reduce costs, members are invited to advise the Membership Secretary (contact details above) of their email address. If any member would like to contact another member by email, they may write to the Membership Secretary and a message will be passed on.

1. Individual Members (by surname in alphabetical order)

- ASHFORD, A., 64 Main Street, Askham Bryan, York, North Yorks. YO23 3QU, **U.K.**
 ASHFORD, R. W., DSc, 142 Meols Parade, Meols CH47 6AN, **U.K.**
- BARGAIN, B., Village de Trunvel, 29720 Treogat, **France**
- BARKER, J.C., Avondale, Westbourne Rd, Westbourne, Sussex PO10 8UL, **U.K.**
- BARLOW, C., c/o D. Hook, 1 Waingate, Grimsargh, Preston, Lancs PR2 5RL, **U.K.**
- BEAMAN, M., Two Jays, Kemple End, Birdy Brow, Stonyhurst, Lancashire BB6 9QY, **U.K.**
- BEECROFT, R., 2 Fen Cottage, Creting St Mary, Ipswich IP6 8QE, **U.K.**
- BELMAN, P.J., 2 School Passage, Southall, London UB1 2DR, **U.K.**
- BERGH, M. VAN DEN, Nieuwezijds Voorburgwal 344, 1012 RX Amsterdam, **The Netherlands**
- BIJLSMA, R., Doldersummerweg 1, 7983 LD, Wapse, **The Netherlands**
- BORROW, N., 18 Knights Green, Sheringham, Norfolk NR26 8UQ, **U.K.**
- BOWDEN, C.G.R., 2 Dutter End, Gamlingay, Sandy, Beds. SG19 3EY, **U.K.**
- BRANSCOMBE, J., Greens, Toab, Orkney KW17 2QG, **U.K.**
- BROADBRIDGE, M.H., 23A Westfield Road, Barton-on-Humber, Humberside DN18 5AA, **U.K.**
- BROUWER, J., Wildekamp 32, 6721 JD Bennekom, **The Netherlands**
- BROWNE, P.W.P., 115 Crichton Street, Ottawa, Ontario K1M 1V8, **Canada**
- BUCKINGHAM, D.L., 1 Lion Mews, Bull St., Potton, Beds. SG19 2NR, **U.K.**

BUIJ, R., Wuitekampweg 5, 6741 CN Lunteren, **The Netherlands**

CHAPPUIS, Dr C., Les Chardonnerets, 10 Vallon du Fer à Cheval, 76530 La Bouille,
France

CHEKE, Prof. R.A., Natural Resources Institute, University of Greenwich at Medway,
Central Avenue, Chatham, Kent ME4 4TB, **U.K.**

CITEGETSE, G., BirdLife International, c/o Conservation Society of Sierra Leone, 4C
Old Railway Line, Tengbeh Town, PO Box 1292, Freetown, **Sierra Leone**

COLLINS, I.D., Bryher, 10 Row Lane, Seend Cleeve, Melksham SN12 6PR, **U.K.**

COOK, A.W., 6 Hazel Avenue, Thame, Oxon. OX9 2AW, **U.K.**

COWAN, DR P., School Croft, Timmerlun Lane, Newburgh AB41 6BF, **U.K.**

DAVIES, M.G., Summerleas, Crapstone Road, Yelverton, Devon PL20 6BT, **U.K.**

DEGAUQUIER, R., 298 rue Jean Jaurès, 59920 Quiévrechain, **France**

DEMETER, A., Av. Van Becelaere 13A, 1170 Brussels, **Belgium**

DEMEY, R., Walter Thijsstraat 9, B-3500 Hasselt, **Belgium**

DODMAN, T., Hundland, Papa Westray, Orkney KW17 2BU, **U.K.**

DOWSETT, R.J., Le Pouget, F-30440 Sumène, **France**

ECCLES, S.D., 8904 Narem Place, Annandale, Virginia 22003, **U.S.A.**

ELLIOT, Sir C.C.H., Bt, Ph.D., Blue Barn House, South Leigh, Witney, Oxon OX29
6XH, **U.K.**

ELLIOTT, A., Sardenya 476 Ent 30, 08025 Barcelona, **Spain**

FANSHAWE, Dr J., Birdlife International, Wellbrook Court, Girton Road, Cambridge
CB3 0NA, **U.K.**

FARNSWORTH, S.J., Hammerkop, Frogmill, Hurley, Maidenhead SL6 5NL, **U.K.**

FIELD, G.D., 37 Milton Grove, New Milton, Hampshire BH25 6HB, **U.K.**

FISHPOOL, Dr L.D.C., 12 Mountain Street, Chilham, Canterbury, Kent CT4 8DQ,
U.K.

FOURIE, S.R., The Orchard, Benthall Lane, Benthall, Broseley, Shrops. TF12 5RR,
U.K.

FRY, Dr C.H., Bridge End House, Kentmere, Staveley, Cumbria LA8 9JP, **U.K.**

FULFORD, Dr T., Medical Research Council, Keneba Field Station, Lower River
Region, **The Gambia**

GALLNER, J.C., St Pierre de Beaujeu, F-04420 La Javie, **France**

GARDINER, S., Ashfoden House, Rodmell, Lewes BN7 3HE, **U.K.**

GATTER, Dr W., Buchstrasse 20, D-73252, Ober Lenningen, **Germany**

GERMAIN, Dr M., 44 rue Cluseret, 92150 Suresnes, **France**

GRAHAM, A.M., Gull's Reach, 11 Armada Drive, Teignmouth TQ14 9NF, **U.K.**

GRANT, A.C., Inverquharity Castle, Kirriemuir, Angus, Scotland, **U.K.**

- GREEN, A.A., 486 Little Mohawk Road, Shelburne Falls, MA 01370, **U.S.A.**
 GRÜNEWALD, Dr J., Stohrerweg 20, D-72070 Tübingen, **Germany**.
 GUITARD, J.-J., "Les ZoiZoZen", Allée Saint-Esprit, 83830 Figanères, **France**
 GULLICK, T., India Alta, Villa Nueva de los Infantes, 13220 Ciudad Real, **Spain**
 GUNNINGHAM, Dr F.J., 5892 Foxtail Court, Ferndale, WA 98248, **U.S.A.**
- HALL, P., OBE [**Nigeria**], A.G. Leventis Ltd, West Africa House, Hanger Lane, Ealing, London W5 3QR, **U.K.**
 HANDKE, C., Stendahlerstr. 13, D-39326 Loitsche, **Germany**
 HARVEY, M.S., 23 Birchfield Lane, Milbarton, Norwich NR14 8BZ, **U.K.**
 HEATON, A., 19 Rydal Gardens, Ashby de la Zouche, Leicestershire LE65 1FJ, **U.K.**
 HIGGLETON, P., 17 Kinlett Close, Highwoods, Colchester CO4 4UE, **U.K.**
 HILLYER, A., 2 Evergreen Drive, 8 Milton Road, Bournemouth, Dorset BH8 8IP, **U.K.**
 HJORT, Dr C., Hessele, Munkarp 111, SE 24391 Hoor, **Sweden**
 HÖLLER, C., Kammakargatan 12, Stockholm 11140, **Sweden**
 HOPKINS, M., TCNN, PO Box 64, Bukuru, Plateau State, **Nigeria**
 HOŘÁK, D., Dept of Ecology, Faculty of Science, Charles University in Prague, Viničná 7, CZ-12844 Praha 2, **Czech Republic**
 HOWE, S., 9 Craighour Avenue, Torphins-by-Banchory, Aberdeen AB31 4JA, **U.K.**
 HOYO, J. DEL, Lynx Edicions, Montseny 8, 08193 Bellaterra (Barcelona), **Spain**
- ISENMANN, P., CEFÉ/CNRS, 1919 route de Mende, F-34293 Montpellier Cedex 5, **France**
- JONES, Dr P.J., Balloan House, Lairg, Sutherland, IV27 4DG, **U.K.**
 JONES, Ms R.M., Flat 1, 16 Vanbrugh Park, Blackheath, London SE3 7AF, **U.K.**
- KELLY, Rev. M., Sonkwala, PO Box 170, Obudu, Cross River State, **Nigeria**
 KING, T., Woodlea House, Stone Drive, Colwall, Malvern, Worcs WR13 6QL, **U.K.**
 KISASA KAFUTSHI, R., 11-8540 Boulevard VIAU, St-Léonard, Québec H1R 2T7, **Canada**
 KLOP, E., M.L. Kingstraat 28, 9728 WJ Groningen, **The Netherlands**
 KOUGOUM, P.G., Dept of Environmental Sciences, Higher Institute of the Sahel, University of Maroua-Cameroon, PO Box 46, Maroua, **Cameroon**.
 KREULEN, D.A., Multatulilaan 58, 9752 GG Haren, **The Netherlands**
 KUHN, I.J., Paul Schneiderstrasse 6, D99423, Weimar, **Germany**
- LACHENAUD, O., Mellier, 79120 Chenay, **France**
 LAESSER, J., Erlenstrasse 17, CH 6020 Emmenbrücke, **Switzerland**
 LAFFARGUE, C., La Tuilerie, 47200 Forques sur Garonne, **France**
 LANG, J.R., 1 Knightwood Court, Rhinefield Rd, Brockenhurst SO41 7UR, **U.K.**
 LANGUY, M., 1 rue de la Bruyère, B-7743 Obigies, **Belgium**
 LEVENTIS, A.P., CBE, 19 Ilchester Place, London W14 8AA, **U.K.**

LIEDÉN, U., Theodorenstr. 13, 65189 Wiesbaden, **Germany**

MANU, Dr S., APLORI [**Nigeria**], c/o P. Hall OBE, A.G. Leventis Ltd, West Africa House, Hanger Lane, Ealing, London W5 3QR, **U.K.**

MANVELL, A., 14 Gloucester Road, Norwich NR2 2DX, **U.K.**

MARTINS, R., 6 Connaught Road, Norwich NR2 3BP, **U.K.**

MASTERTSON, A., Cottage 3, B.S. Leon, Box A 80 Avondale, **Zimbabwe**

MILLS, T.R., Cruglas Swyddffynnon, Ystrad Meurig, Ceredigion SY25 6AN, **U.K.**

MOORE, A.M., 1 Uppingham Road, Oakham, Rutland LE15 6JB, **U.K.**

MOREL, Dr M.-Y., 43 rue Fessart, 92100 Boulogne-Billancourt, **France**

MUNDY, P., PO Box FM 424, Famona, Bulawayo, **Zimbabwe**

NASASAGARE, R.P., Ecole Normale Supérieure, BP 6983 Bujumbura, **Burundi**

NELSON, M., 1629 El Prado Dr., Knoxville, TN 37922, **U.S.A.**

NESSELROADE, D., 738 Linmuth Ct., Springfield, OH 45503, **U.S.A.**

OTTOSSON, Dr U., 18A Rue de Mamer, L-8280 Kehlen, **Luxembourg**

PARK, P.O., 45 Hills Road, Cambridge CB2 1NZ, **U.K.**

PASSAVY, G., 25 rue Erard, 75012 Paris, **France**

PAYNE, Dr R.B., 1306 Granger Ave, Ann Arbor, MI 48104, **U.S.A.**

PHALAN, B., Zoology Dept, Downing St, Cambridge CB2 3EJ, **U.K.**

PORTIER, B., Chemin de la Neuville 40, 4821 Andrimont, **Belgium**

QUANTRILL, W., Tor House, 36 Newtown, Bradford-on-Avon, Wiltshire BA15 1NF, **U.K.**

RAINEY, H., 140 Argyle St, Cambridge CB1 3LS, **U.K.**

RANDALL, G., 6 Abbeyfields, Abbey Rd, Great Massingham, King's Lynn, Norfolk PE32 2JE, **U.K.**

REDMAN, N., 36 Soho Square, London W1D 3QZ, **U.K.**

ROBB, M., 2300 Rue Grenet, Apt #311, St Laurent, Québec H4L 4Y9, **Canada**

ROBIN, N., 35 Rue Bonaparte, 75006 Paris, **France**

ROSE, J., 1 bis Rue des Châtre-Sacs, 92310 Sèvres, **France**

RUMSEY, S., Elms Farm, Pett Lane, Icklesham, Winchelsea, E Sussex TN36 4AH, **U.K.**

SALEWSKI, V., Prinz-Rupprechtstr 24, 93053 Regensburg, **Germany**

SCHOLTE, P.T., Nieuwe Teentuin 12c, 1013 LV Amsterdam, **The Netherlands**

SHORT, Dr L., PMB, Nanyuki, **Kenya**

SIKKEMA, M., Eilansgrien 36, 9264 TC Earnewald, **The Netherlands**

SINEUX, L. [**Ivory Coast**], Joris van der Haagenlaan 9, 6814 LJ Arnhem, **The Netherlands**

SKILLETER, M., 12 Westcliffe Road, Roker, Sunderland, **U.K.**

SKINNER, Dr N.J., 48 Everitt Court, Swonells Walk, Oulton Broad, Lowestoft NR32 3PL, **U.K.**

SMITH, V.W., 1 Karrakatta Road, Goode Beach, Albany, Western Australia 6330, **Australia**

TAMUNGANG, A.S., Dept of Animal Biology, University of Dschang, PO Box 146, Dschang, West Province, **Cameroon**

THIOLLAY, Dr J.-M., 2 rue Rivière, F-10220 Rouilly Sacey, **France**

THOMA, M., Gesellschaftsstr. 89, 3012 Bern, **Switzerland**

THOMPSON, Dr H.S., 6 Topham Way, Cambridge, Cambs CB4 3RA, **U.K.**

TURNER, D.A., PO Box 1651, 20117 Naivasha, **Kenya**

TYE, Dr A., PO Box 31, Kalo Chorio Oreinis, Nicosia 2616, **Cyprus**

VOADEN, N.J., 18 Fair Hill, Shipham, Winscombe, Somerset BS2 1TH, **U.K.**

WACHER, Dr T., Conservation Programmes, Zoological Society of London, Regents Park, London NW1 4RY, **U.K.**

WAGNER HODGES, Ms V.J., 12 North View, Wimbledon Common, London SW19 4UJ, **U.K.**

WALL, J.W., 19 Tisdale Road, Scarsdale, New York, NY 10583-5613, **U.S.A.**

WALLACE, J.P., 50 Cherryburn Gardens, Fenham, Newcastle-upon-Tyne NE4 9UO, **U.K.**

WALSH, J.F., 80 Arundel Road, Lytham St Annes, Lancs. FY8 1BN, **U.K.**

WALTERT, Dr M., Georg-August-Universität Göttingen, Dept of Conservation Biology, Bürgerstrasse 50, 37073 Göttingen, **Germany**

WILKINSON, Dr R., North of England Zoological Society, Zoological Gardens, Cheshire CH2 1LH, **U.K.** <r.wilkinson@chesterzoo.co.uk>

WOODBRIDGE, K., Twinsness, North Ronaldsay, Orkney KW17 2BE, **U.K.**

WOODCOCK, M., Furlongs, Blakeney Long Lane, Wiveton, Norfolk NR25 7DD, **U.K.**

WÜST, R., Faberstr.4, D-70188 Stuttgart, **Germany**

2. Institutional Members

(arranged in alphabetical order of continent, then country)

Africa

CAMEROON BIODIVERSITY CONSERVATION SOCIETY, c/o WCS, B.P. 3055, Messa, Yaoundé, **Cameroon**

Bibliothécaire, Faculté des Sciences, UNIVERSITÉ DE YAOUNDÉ, **Cameroon**

N.C.R.C., PO Box 925, Kasheshie, Accra, **Ghana**

Bibliothécaire, Laboratoire de Zoologie, Faculté des Sciences, UNIVERSITÉ DE
COCODY, 22 BP582, Abidjan 22, **Ivory Coast**

BIRDLIFE AFRICA, ICIPE Campus, Kasarani Road, POB 3502-00100, **Kenya**

Librarian, NATUREKENYA, National Museums of Kenya, PO Box 40658, Nairobi,

Kenya

Librarian, Kashim Ibrahim Library, AHMADU BELLO UNIVERSITY, Zaria, **Nigeria**

A.P. LEVENTIS ORNITHOLOGICAL INSTITUTE, Fobel Rd, Laminga, Jos East, LGA Jos,

Nigeria

Bibliothécaire, DIRECTION DES PARCS NATIONAUX DU SÉNÉGAL, BP 5135, Dakar,

Sénégal

Bibliothécaire, STATION D'ORNITHOLOGIE, I.R.D., BP 1386, Dakar, **Sénégal**

Bibliothécaire, Faculté des Sciences, UNIVERSITÉ DE DAKAR, Dakar, **Sénégal**

Librarian, DURBAN NATURAL HISTORY MUSEUM, PO Box 4085, Durban 4000, **South**

Africa

Margaret Koopman, Niven Library, PERCY FITZPATRICK INSTITUTE, University of

Cape Town, Rondebosch, Western Cape 7701, **South Africa**

BIRDLIFE ZIMBABWE, PO Box RV100, Runiville, Harare, **Zimbabwe**

Librarian, NATIONAL MUSEUM, PO Box 240, Bulawayo, **Zimbabwe**

Europe

ROYAL MUSEUM OF CENTRAL AFRICA, Central Library, Leuvensesteenweg 13, 3080

Tervuren, **Belgium**

ZOOLOGICAL MUSEUM Library, Universitetsparken 15, 2100 København Ø, **Denmark**

MUSÉUM NATIONAL D'HISTOIRE NATURELLE, Bibliothèque des Mammifères et des

Oiseaux, 57 rue Cuvier, 75231 Paris Cedex 05, **France**

OFFICE NATIONAL DE LA CHASSE, CNER Avifaune Migratrice Station, 85340 L'Ile

d'Olonne, **France**

Bibliothécaire, STATION BIOLOGIQUE DE LA TOUR DU VALAT, Fondation Sansouire Le

Sambuc, F-13200 Arles, **France**

INSTITUT FÜR VOGELFORSCHUNG, Vogelwarte Helgoland, 26386 Wilhelmshaven,

Rustereil, **Germany**

MUSEUM KOENIG, The Library, Adenauerallee 160, D-53113 Bonn, **Germany**

UNIVERSITÄTSBIBLIOTHEK JOHANN CHRISTIAN SENCKENBERG, Zietschriftenabteilung/DFG,

Bockenheimerlandstraße 134–138, D-60325 Frankfurt am Main, **Germany**

The Librarian, A&W ECOLOGICAL CONSULTANTS, POB 32, 9269 ZR Feanwâlden, **The**

Netherlands

The Librarian, NATIONAAL NATUURHISTORISCH MUSEUM NATURALIS, Bibliotheek

Naturalis, Postbus 9517, 2300 RA Leiden, **The Netherlands**

The Librarian, VOGELBESCHERMING/BIRDLIFE-THE NETHERLANDS, Postbus 925, NL-

3700 AX Zeist, **The Netherlands**

MUSEUM AND INSTITUTE OF ZOOLOGY, Polish Academy of Science, Library, Wilcza St

64, PL-00-679 Warszawa, **Poland**

THE RING, (Prof L. Tomiałojc), Natural History Museum, Siendiewicza 21, 50335
Wroclaw, **Poland**

BIRDLIFE SLOVENIA, PP 2990, SI-1001 Ljubljana, **Slovenia**

LUNDS UNIVERSITET, Biologibiblioteket, Solvegatan 237, SE-223 62 Lund, **Sweden**

Library, SWISS ORNITHOLOGICAL INSTITUTE, 6204 Sempach, **Switzerland**

AFRICAN BIRD CLUB, The Secretary, c/o BirdLife International, Wellbrook Court,
Girton Road, Cambridge CB3 0NA, **U.K.**

AGENCY FOR THE LEGAL DEPOSIT LIBRARIES, Causewayside Building, 33 Salisbury
Place, Edinburgh EH9 1SL, **U.K.**

The Librarian, BIRDLIFE INTERNATIONAL, Wellbrook Court, Girton Road, Cambridge
CB3 0NA, **U.K.**

BRITISH LIBRARY Acquisitions Unit (SR15), Boston Spa, Wetherby LS23 7BQ, **U.K.**

Librarian, Dept of Zoology, EDWARD GREY INSTITUTE, South Parks Road, Oxford
OX1 3PS, **U.K.**

NATURAL HISTORY MUSEUM — Acquisitions Section, Library & Archives, Cromwell
Rd, London SW7 5BD, **U.K.**

ZOOLOGICAL RECORD (Thomson Reuters), Enterprise House, Innovation Way,
Heslington, York YO10 5NY, **U.K.**

North America

Librarian, ROYAL ONTARIO MUSEUM, University of Toronto, 100 Queens Park,
Toronto ON, M5S 2C6, **Canada**

Library Serials Unit/P. Watson, AMERICAN MUSEUM OF NATURAL HISTORY, Central
Park West at 79th Street, New York, NY 10024-5192, **U.S.A.**

CORNELL UNIVERSITY LIBRARY, E-resources & Serials Management, 110 Olin
Library, Ithaca, NY 14853, **U.S.A.**

FIELD MUSEUM OF NATURAL HISTORY, Library Subs, 1400 Lake Shore Drive, Chicago
IL 60605-2498, **U.S.A.**

Ernst Mayr Library, MUSEUM OF COMPARATIVE ZOOLOGY, Harvard University, 26
Oxford St, Cambridge, MA 02138, **U.S.A.**

Ornithology Library, PEABODY MUSEUM OF NATURAL HISTORY, Yale University, PO
Box 208118, New Haven, Connecticut 06520, **U.S.A.**

SMITHSONIAN INSTITUTION Libraries, NHB 25 MRC 154, PO Box 37012, Washington,
DC 20013-7012, **U.S.A.**

WILSON ORNITHOLOGICAL SOCIETY, Museum of Zoology, University of Michigan,
Ann Arbor, Michigan 48104, **U.S.A.**

UNIVERSITY OF WISCONSIN, Memorial Library, Acquisition Serials Dept, 728 State
Street, Madison, WI 53706-1418, **U.S.A.**

Instructions aux Auteurs

Malimbus publie des articles de recherche, des revues de publications et des nouvelles traitant de l'ornithologie ouest-africaine.

Les **Articles** et les **Notes Courtes** doivent être des apports originaux; ceux déjà publiés ailleurs, en partie ou en totalité, seront normalement refusés. Les Notes Courtes sont des articles de moins de 1500 mots (références comprises) ou de quatre pages imprimées. Autant que possible, les manuscrits auront été au préalable soumis à au moins un ornithologue ou biologiste pour un examen minutieux. Les manuscrits seront envoyés pour critique à au moins un lecteur compétent.

Les textes des **Nouvelles & Commentaires** ne devraient pas dépasser 1000 mots.

Les **textes** sont acceptés en anglais et en français; la Rédaction pourra aider les auteurs dont la langue maternelle n'est pas l'une de celles-ci. Nous préférons les envois de manuscrits par email (en pièce jointe). Consultez le Rédacteur pour plus de détails, par ex. les logiciels compatibles.

Tous les Articles (mais non les Notes Courtes) comporteront un **Résumé**, n'excédant pas 5% de la longueur totale. Le Résumé mentionnera brièvement les principaux résultats et conclusions de l'Article et ne sera pas un simple compte rendu de ce qui a été fait. Les résumés seront publiés à la fois en anglais et en français (ou dans la langue officielle du pays dans lequel le travail a été réalisé) et seront traduits au mieux par la Rédaction.

La **présentation** des tableaux, chiffres, unités métriques, références, *etc.* doit correspondre à celles des numéros récents. A notez, en particulier: Les noms des auteurs doivent être listés en faisant précéder le nom de famille par le prénom ou des initiales (ex. John A. Smith); les dates seront écrites "2 fév 1990" mais les mois seuls pourront être écrits en entier; les heures seront écrites "6h45", "17h00"; les coordonnées "7°46'13"N" (pas de zéros en tête) ou en degrés décimaux jusqu'à cinq décimales (ex. 1.23456°N), mais non en minutes décimales; les nombres jusqu'à dix seront écrits en toutes lettres, excepté devant une unité de mesure (ex. 6 m); les nombres à partir de 11 seront écrits en chiffres sauf au début d'une phrase. Toutes les références citées dans l'article, et aucune autre, doivent figurer dans la bibliographie.

Les **articles sur l'avifaune** doivent comprendre une carte ou un index géographique, incluant tous les endroits cités. Ils doivent comporter quelques brèves indications sur le climat, la topographie, la végétation et les circonstances ou événements inhabituels avant ou pendant l'étude (ex. pluies tardives, *etc.*). Les **listes d'espèces** ne doivent contenir que des données importantes: les listes complètes ne sont justifiées que pour les régions encore non étudiées ou délaissées pendant longtemps. Autrement, ne citer que les espèces sur lesquelles l'étude fournit une information nouvelle sur la répartition, la période de séjour, la reproduction, *etc.* Pour chaque espèce, indiquer l'extension de l'aire de répartition, une estimation d'abondance (*Malimbus* 17: 38) et les données datées sur la reproduction; indiquer le statut migratoire et la période de séjour seulement tels qu'ils ressortent de l'étude. Eventuellement, remplacer les données dans le contexte en les comparant brièvement avec une liste régionale de référence. Les longues listes d'espèces peuvent être présentées sous la forme de tableaux (ex. *Malimbus* 25: 4–30, 24: 15–22, 23: 1–22, 1: 22–28, or 1: 49–54) ou sous la forme rédigée des numéros récents. La **séquence taxonomique** et les **noms scientifiques** (et de préférence aussi les **noms vernaculaires**) doivent suivre Borrow & Demey (2001, *Birds of Western Africa*, Christopher Helm, London, avec des noms rectifiés de Borrow & Demey 2004, *Field Guide to the Birds of Western Africa*, Christopher Helm, London), ou *The Birds of Africa* (Brown *et al.* 1982, Urban *et al.* 1986, Fry *et al.* 1988, Keith *et al.* 1992, Fry & Keith 2000, 2004, Academic Press, London), à moins de donner les raisons de s'écarter de ces auteurs. Un **guide plus complet à l'intention aux auteurs** d'articles sur l'avifaune, comprenant l'échelle d'abondance des espèces conseillée, a été publié dans *Malimbus* 17: 35–39 et une version augmentée et actualisée de celle-ci mise sur le site internet (<http://malimbus.free.fr/instmale.htm>). La Rédaction se fera un plaisir de donner des conseils pour les études spécifiques.

Pour le dessin des **Figures**, et en particulier la taille des caractères, tenir compte des dimensions de la page de *Malimbus*. On préfère les figures préparées sur logiciel graphique approprié et sauvegardées en haute définition. Les fichiers de basse résolution et les impressions de mauvaise qualité seront refusés. Les auteurs sont encouragés à soumettre des **photographies** qui illustrent des points importants de leurs articles. Les photographies doivent être en couleurs et de haute définition. Les figures et les photographies doivent être envoyées comme fichiers de logiciel graphique (par ex. jpg ou tif), et non pas être incluses dans un fichier de Word. Consulter le Rédacteur pour tout renseignement.

Un fichier pdf des Articles et des Notes Courtes, et une copie du numéro de publication seront envoyés gratuits à l'auteur ou à l'auteur principal.



MALIMBUS 36(2) September 2014

Contents — Table des Matières

| | |
|---|---------|
| Editorial: Our 50th Anniversary | 65–66 |
| Raptors in Yankari Game Reserve and surrounding unprotected area, Nigeria. | |
| J.D. Onoja, T. Tende, T.C. Omotoriogun, U. Ottosson, S.A. Manu & G.S. Mwansat | 67–75 |
| Birds of the International Institute of Tropical Agriculture campus, a stronghold of avian diversity in the changing Ibadan area (Nigeria) over the last 50 years. | |
| T.A. Adeyanju, U. Ottosson, T.E. Adeyanju, T.C. Omotoriogun, P. Hall, S.A. Manu, T. Alabi, G.A. Lameed & D. Bown | 76–105 |
| L'avifaune d'un milieu de riziculture et de ses environs dans la zone humide de Grand-Bassam, sud-est Côte d'Ivoire. | |
| G.S.K. Oudouké, H.K. Yaokokoré-Béibro, M.E. Konan & P.K. Kouadio | 106–115 |
| Short Notes — Notes Courtes | |
| Records of African Pygmy Goose <i>Nettapus auritus</i> and a <i>Haleyon</i> kingfisher on São Tomé. | |
| B. Hansson & A. Thomasson | 116–117 |
| Observations ornithologiques au Sénégal et en Gambie. | |
| M.S. Diop, B. Mennesson & J. Rose | 117–119 |
| The Parelus bird collection from Ivory Coast at the Field Museum of Natural History, and the first country record of Rufous <i>Cisticola Cisticola rufus</i>. | |
| J.I. Engel, D. Parelus & J.M. Bates | 119–122 |
| News | 123 |
| Society Notices — Informations de la Société | 124 |